

Evidence-Based Guidelines for Weaning and Discontinuing Ventilatory Support

A Collective Task Force Facilitated by the American College of Chest Physicians, the American Association for Respiratory Care, and the American College of Critical Care Medicine

Introduction

Pathophysiology of Ventilator Dependence

Criteria to Assess Ventilator Dependence

Managing the Patient Who Has Failed a Spontaneous Breathing Test

Role of Tracheotomy in Ventilator-Dependent Patients

The Role of Long-Term Facilities

[Respir Care 2002;47(1):69–90]

Introduction

The discontinuation or withdrawal process from mechanical ventilation is an important clinical issue.^{1,2} Patients are generally intubated and placed on mechanical ventilators when their own ventilatory and/or gas exchange capabilities are outstripped by the demands placed on them from a variety of diseases. Mechanical ventilation also is required when the respiratory drive is incapable of initiating ventilatory activity, either because of disease processes or drugs. As the conditions that warranted placing the patient on the ventilator stabilize and begin to resolve, attention should be placed on removing the ventilator as

quickly as possible. Although this process often is termed “ventilator weaning” (implying a gradual process), we prefer the more encompassing term “discontinuation.”

SEE THE RELATED EDITORIAL ON PAGE 29

Unnecessary delays in this discontinuation process increase the complication rate from mechanical ventilation (eg, pneumonia, airway trauma) as well as the cost. Aggressiveness in removing the ventilator, however, must be balanced against the possibility that premature discontinuation may occur. Premature discontinuation carries its own set of problems, including difficulty in reestablishing artificial airways and compromised gas exchange. It has been estimated that as much as 42% of the time that a medical patient spends on a mechanical ventilator is during the discontinuation process.³ This percent is likely to be much higher in patients with more slowly resolving lung disease processes.

There are a number of important issues involved in the management of a mechanically ventilated patient whose disease process has begun to stabilize and/or reverse such that the discontinuation of mechanical ventilation becomes a consideration. First, an understanding of all the reasons that a given patient required a mechanical ventilator is needed. Only with this understanding can medical management be optimized. Second, assessment techniques to identify patients who are capable of ventilator discontinuation need to be utilized. Ideal assessment techniques should be able to easily and safely distinguish which pa-

The Writing Committee, on behalf of the Panel, includes Neil R MacIntyre MD FAARC (Chairman), Deborah J Cook MD, E Wesley Ely Jr MD MPH, Scott K Epstein MD, James B Fink MSc RRT FAARC, John E Heffner MD, Dean R Hess PhD RRT FAARC, Rolf D Hubmayr MD, and David J Scheinhorn MD. Other members of the panel include Suzanne Burns RN MSN CCRN RRN, David Chao MD, Andres Esteban MD, Douglas R Gracey MD, Jesse B Hall MD, Edward F Haponik MD, Marin H Kollef MD, Jordi Mancebo MD, Constantine A Manthous MD, Arthur S Slutsky MD, Meg A Stearn-Hassenpflug MS RD, and James K Stoller MD MSc FAARC.

Reprinted with permission from *Chest* (Chest 2001;120(6)375–395). © American College of Chest Physicians. Copies can be ordered from the American College of Chest Physicians, at 1-800-343-2227 or 1-847-498-1400. www.chestjournal.org.

Correspondence: Neil R MacIntyre MD FAARC, Respiratory Care Services, Duke University Medical Center, Box 3911, Durham NC 27710. E-mail: neil.macintyre@duke.edu.

Table 1. Grades of Evidence

Grade	Description
A	Scientific evidence provided by well-designed, well-conducted, controlled trials (randomized and nonrandomized) with statistically significant results that consistently support the guideline recommendation
B	Scientific evidence provided by observational studies or by controlled trials with less consistent results to support the guideline recommendation
C	Expert opinion supported the guideline recommendation, but scientific evidence either provided inconsistent results or was lacking

tients need prompt discontinuation and which need continued ventilatory support. Third, ventilator management strategies for stable/recovering patients who still require some level of ventilatory support need to be employed. These strategies need to minimize both complications and resource consumption. Fourth, extended management plans (including tracheotomy and long-term ventilator facilities) need to be considered for the long-term ventilator-dependent patient.

To address many of these issues, the Agency for Healthcare Policy and Research (AHCPR) charged the McMaster University Evidence Based Practice Center to do a comprehensive evidence-based review of many of the issues involved in ventilator weaning/discontinuation. Led by Deborah Cook MD, an exhaustive review of several thousand articles in the world literature resulted in a comprehensive assessment of the state of the literature in 1999.⁴ At the same time, the American College of Chest Physicians (ACCP), the Society for Critical Care Medicine (SCCM), and the American Association for Respiratory Care (AARC) formed a task force to produce evidence-based clinical practice guidelines for managing the ventilator-dependent patient during the discontinuation process. The charge of this task force was to utilize the McMaster AHCPR report as well as their own literature review to address the following 5 issues: (1) the pathophysiology of ventilator dependence; (2) the criteria for identifying patients who are capable of ventilator discontinuation; (3) ventilator management strategies to maximize discontinuation potential; (4) the role of tracheotomy; and (5) the role of long-term facilities. Review/writing teams were formed for each of these issues.

From these evidence-based reviews, a series of recommendations were developed by the task force, which are the basis of this report. Each recommendation is followed by a review of the supporting evidence, including an assessment of the strength of the evidence (Table 1). As there were many areas in which evidence was weak or absent, the expert opinion of the task force

was relied on to “fill in the gaps.” Consensus was reached, first, by team discussions and, later, through the repeated cycling of the draft through all members of the task force.

Both the McMaster AHCPR group and the task force recognized the needs for the future. These include more randomized controlled trials to look at a number of issues. Among the more important questions that need answering are the following: (1) Which criteria are the best indicators of reversal of respiratory failure in the screening process? (2) What factors are involved in ventilator dependence and which measurement techniques are most useful in determining ultimate success in the discontinuation process? (3) In balancing discontinuation aggressiveness against the risks of premature discontinuation, what is a reasonable reintubation rate in patients recently removed from ventilatory support? (4) What is the value of trying to reduce levels of partial ventilator support in stable/recovering patients who have failed a discontinuation assessment? (5) What role do tracheotomies have in facilitating the discontinuation process? (6) What is the role of the long-term facility, and when should patients be transferred to such facilities?

Pathophysiology of Ventilator Dependence

Introduction

Patients require mechanical ventilatory support when the ventilatory and/or gas exchange capabilities of their respiratory system fail. This failure can be the result of processes both within the lung as well as in other organ systems, most notably the central nervous and the cardiovascular systems. Although patients may be dependent on ventilatory support for brief periods of anesthesia or neuromuscular blockade, the term “ventilator-dependent” is usually reserved for patients with a need for mechanical ventilation beyond 24 hours or by the fact that they have failed to respond during discontinuation attempts. Under these circumstances, the clinical focus should be not only on ventilator management but also should include a search for all of the possible reasons (especially potentially reversible ones) that may explain the ventilator dependency.

Recommendation 1. *In patients requiring mechanical ventilation for > 24 hours, a search for all the causes that may be contributing to ventilator dependence should be undertaken. This is particularly true in the patient who has failed attempts at withdrawing the mechanical ventilator. Reversing all possible ventilatory and nonventilatory issues should be an integral part of the ventilator discontinuation process.*

Table 2. Causes of Ventilator Dependency

Causes	Description
Neurologic controller	Central drive; peripheral nerves
Respiratory system	Mechanical loads: respiratory system mechanics; imposed loading Ventilatory muscle properties: inherent strength/endurance; metabolic state/nutrients/oxygen delivery and extraction Gas exchange properties: vascular properties and ventilation/perfusion matching
Cardiovascular system	Cardiac tolerance of ventilatory muscle work; peripheral oxygen demands
Psychological issues	

Evidence (Grade B)

There are a number of specific reasons why patients may be ventilator-dependent (Table 2). Determining which factor or factors may be involved in a given patient requires both clinical awareness of these factors as well as focused clinical assessments. The search for the underlying causes for ventilator dependence may be especially important if previously unrecognized, but reversible, conditions are discovered.

Neurologic Issues: The ventilatory pump controller in the brainstem is a rhythm and pattern generator, which receives feedback from cortical, chemoreceptive and mechanoreceptive sensors. The failure of this controller can come from several factors.⁵⁻¹² These factors can be either structural (eg, brainstem strokes or central apneas) or metabolic (eg, electrolyte disturbances or sedation/narcotic usage^{13,14}). The failure of the peripheral nerves also can be the result of either structural factors¹⁵ or metabolic/drug factors.^{16,17} A unique neurologic dysfunction that also could cause ventilator dependence is obstructive sleep apnea, in which an artificial airway may be necessary to maintain airway patency.^{10,11}

Respiratory System Muscle/Load Interactions: Often, patients who exhibit ventilator dependence do so because there appears to be a mismatch between the performance capacity of ventilatory pump and the load placed on it (ie, the capacity/load imbalance hypothesis).¹⁸⁻²³ There is ample evidence that ventilatory pump performance may be impaired in ventilator-dependent patients because ventilatory muscles are weak. This may be a consequence of atrophy and remodeling from inactivity.^{2,24} It may also be a consequence of injury from overuse and of insults associated with critical illness neuropathy and myopathy.²⁵⁻²⁹ A number of drugs (eg, neuromuscular blockers, aminoglycosides, and corticosteroids) also can contribute to myopathy,^{17,30-32} as can various metabolic derangements (see

below). Finally, dynamic hyperinflation can put ventilatory muscles in a mechanically disadvantageous position.³³ In a number of studies, patients who failed to respond to a withdrawal from mechanical ventilation tended to be weaker (ie, they had a lower performance capacity) than those who succeeded,³⁴⁻⁴⁹ but, in general, the within-group variability in respiratory muscle strength was too large to justify general conclusions.

Ventilatory muscle fatigue also could contribute to poor muscle performance. However, the role of fatigue in ventilator dependence is not well understood, and the studies performed to date^{21,26,50-54} have failed to delineate the sensitivity and specificity of specific fatigue tests in ventilator-dependent patients. Ventilatory support reduction-related changes in transdiaphragmatic pressure, respiratory rate, and thoracoabdominal dyssynchrony are clearly not specific manifestations of respiratory muscle fatigue.⁵⁵⁻⁶⁰ The most promising diagnostic test of diaphragm contractility to date is the transdiaphragmatic pressure measurement during twitch stimulation of the phrenic nerves.^{21,61} However, too few patients have been studied with this technique to draw any meaningful conclusions about the prevalence of diaphragm fatigue that is attributable to ventilator dependence.

The load on the ventilatory muscles is a function of ventilation demands and respiratory system mechanics (ie, primarily compliance and resistance). Normal minute ventilation during spontaneous breathing is generally < 10 L/min, normal respiratory system compliance (ie, tidal volume/static inflation pressure) is generally > 50-100 mL/cm H₂O, and normal airway resistance (ie, peak-static inflation pressure/constant inspiratory flow) is generally < 5-15 cm H₂O/L/s. Ventilation demands can increase as a consequence of increased oxygen demands in patients with sepsis or increased dead space in patients with obstructive diseases. Compliance worsening can be a consequence of lung edema, infection, inflammation, or fibrosis, and of chest wall abnormalities such as edema or surgical dressings. Resistance worsening can be a consequence of bronchoconstriction and airway inflammation. Additional load can also be imposed by narrow endotracheal tubes and by insensitive or poorly responsive ventilator demand valves.

The load imposed by ventilation demands interacting with respiratory system mechanics can be expressed as respiratory work, pressure time integral, or the change in metabolism (eg, the oxygen cost attributable to breathing). Many studies^{19,35,62-66} show that patients who are ventilator-dependent tend to have larger respiratory muscle loads than do patients who can be withdrawn from mechanical ventilation. In patients with airways obstruction, the load imposed by dynamic hyperinflation has received particular attention as an important contributor to ventilator depen-

dence.^{23,33,65,67-71} As is true for measures of ventilatory pump capacity, however, most investigators report considerable overlap in load parameters between patients with different discontinuation outcomes.

Patients who go on to fail to respond to ventilator withdrawal attempts because of a capacity/load imbalance tend to display rapid, shallow breathing patterns.^{2,72,73} This pattern is advantageous from an energetics perspective, but it is also associated with increased dead space and wasted ventilation and, hence with impaired carbon dioxide elimination. Chemoreceptive and mechanoreceptive feedback into the neural control of breathing is not well understood, and thus it is difficult to distinguish whether this breathing pattern is a consequence of a reduced respiratory drive per breath or an inability of ventilatory muscles to respond to an appropriately increased neural stimulus.^{19,62,65,71,72}

Metabolic Factors and Ventilatory Muscle Function: Nutrition, electrolytes, hormones, and oxygen transport are all metabolic factors that can affect ventilatory muscle function. Inadequate nutrition leads to protein catabolism and loss of muscle performance.^{74,75} The normal hypoxic ventilatory response and the hypercapnic ventilatory response also have been shown to deteriorate under conditions of semistarvation.⁷⁶ In contrast, overfeeding also can impair the ventilator withdrawal process by leading to excess carbon dioxide production, which can further increase the ventilation loads on ventilatory muscles. Studies^{77,78} have suggested that proper nutritional support can increase the likelihood of success of ventilator withdrawal. A number of electrolyte imbalances also can impair ventilatory muscle function.^{5,9,79-81} Phosphate deficiency has been associated with respiratory muscle weakness and ventilator withdrawal failure. A study demonstrating improved transdiaphragmatic pressure values with the repletion of serum phosphorus levels in patients receiving mechanical ventilation, however, did not specifically address the issue of ventilator withdrawal.⁷⁹ Magnesium deficiency also has been reported to be associated with muscle weakness,⁸² although the relationship to ventilator dependence has not been specifically addressed. Finally, bicarbonate excretion from inappropriate overventilation (often occurring in patients with chronic obstructive pulmonary disease (COPD) with chronic baseline hypercapnia) can impair ventilator withdrawal efforts as the patient has a diminished capacity to compensate for hypercapnia. Severe hypothyroidism and myxedema directly impair diaphragmatic function and blunt ventilatory responses to hypercapnia and hypoxia.^{83,84} Other hormonal factors that are important for optimal muscle function include insulin/glucagon and adrenal corticosteroids.

As in other organs, adequate oxygen delivery and oxygen uptake by the ventilatory muscles is necessary for

proper muscle function.^{85,86} Impaired oxygen delivery can be a consequence either of inadequate oxygen content or of inadequate cardiac output.⁸⁷ Impaired oxygen uptake occurs most commonly during systemic inflammatory syndromes such as sepsis.⁸⁸

Gas Exchange Factors: Gas exchange abnormalities can develop during ventilatory support reductions for several reasons. Various lung diseases produce ventilation-perfusion imbalances and shunts. Ventilator dependence thus may be a consequence of a need for high levels of expiratory pressure and/or the fraction of inspired oxygen ($F_{I_{O_2}}$) to maintain an adequate oxygen content.^{5,9} A patient with hypoxemia also can develop a fall in mixed venous P_{O_2} levels from the cardiovascular factors described below.

Cardiovascular Factors: Several groups of investigators have drawn attention to cardiovascular responses in ventilator-dependent patients and have emphasized the potential for ventilatory support reductions to induce ischemia or heart failure in susceptible patients with limited cardiac reserve.^{9,89-93} Putative mechanisms include the following: (1) increased metabolic demand and hence circulatory demands that are associated with the transition from mechanical ventilation to spontaneous breathing in patients with limited cardiac reserve; (2) increases in venous return as the contracting diaphragm displaces blood from the abdomen to the thorax; and (3) the increased left ventricular afterload that is imposed by negative pleural pressure swings. Lemaire and colleagues demonstrated left ventricular dysfunction (ie, the pulmonary capillary wedge pressure increased from 8 to 25 mm Hg) during failed ventilator withdrawal attempts in 15 patients with COPD. Following diuresis, 9 of these 15 patients were successfully withdrawn from the ventilator.⁹⁰

Psychological Factors: Psychological factors may be among the most important nonrespiratory factors leading to ventilator dependence. Fear of the loss of an apparent life support system as well as social/familial/economic issues all may play a role. Stress can be minimized by frequent communication among the staff, the patient, and the patient's family.⁹⁴ Environmental stimulation using television, radio, or books also appears to improve psychological functioning.² Ambulation using a portable ventilator (or bagging) has been shown to benefit attitudes and outlooks in long-term ventilator-dependent patients. Sleep deprivation may cause impairment of the respiratory control system,⁹⁵ although this may be related to accompanying factors rather than to sleep deprivation per se.⁹⁶ Finally, biofeedback may be helpful in decreasing the weaning time in patients who are having difficulty withdrawing from ventilatory support.⁹⁷⁻⁹⁸

Criteria to Assess Ventilator Dependence

Introduction

The process of discontinuing mechanical ventilatory support begins with a recognition of adequate recovery from acute respiratory failure. Thereafter, careful clinical assessments are required to determine the patient's readiness for subsequent discontinuation of ventilatory support and, ultimately, extubation. To facilitate this process, investigators have focused on identifying objective criteria to determine the answers to the following questions: When can efforts to discontinue ventilation be initiated? What assessment strategies will best identify the patient who is ready for ventilator discontinuation? When should extubation be carried out, and how can extubation outcome best be predicted?

Evidence to answer these questions comes largely from observational studies in which a certain parameter (or set of parameters) is compared in a group of patients who either successfully or unsuccessfully have been removed from the ventilator. The general goal of these studies is to find "predictors" of outcome. Evaluating the results from these types of studies can be difficult for several reasons.

First, the "aggressiveness" of the clinician/investigator's weaning and discontinuation philosophy needs to be understood, as it will affect the performance of a given predictor. A very aggressive clinical philosophy will maximize the number of patients withdrawn from ventilatory support but could also result in a number of premature discontinuations, with a subsequent need for reintubations and/or reinstatement of support. In contrast, a less aggressive clinical philosophy will minimize premature discontinuations but could also unnecessarily prolong ventilatory support in other patients. Unfortunately, there are no good data to help clinicians to determine the best balance between premature and delayed discontinuations in evaluating a given discontinuation strategy. Clearly, extubation failure should be avoided whenever possible because the need for reintubation carries an 8-fold higher odds ratio for nosocomial pneumonia⁹⁹ and a 6-fold to 12-fold increased mortality risk.^{100–103} In contrast, the maintenance of unnecessary ventilator support carries its own burden of patient risk for infection and other complications.^{104,105} Reported reintubation rates range from 4% to 23% for different intensive care unit (ICU) populations,^{100,101,103,104,106–110} and may be as high as 33% in patients with mental status changes and neurological impairment.¹⁰³ Although the optimal rate of reintubation is not known, it would seem likely to rest between 5% and 15%.

Second, a number of methodological problems exist with most of these observational studies. For instance, patients are recruited into these studies because investigators believe that there is some reasonable chance of success for

ventilator discontinuation. These "entry" criteria often include some form of clinical judgment or intuition, making results from one study difficult to compare to another. In addition, clinician/investigators deciding to proceed with ventilator discontinuation/extubation often have not been blinded to the parameters being analyzed as possible predictors. Indeed, the parameter being analyzed may often enter into the clinical decision on whether either to continue or to discontinue ventilatory support. Other methodological problems with these observational studies include different measurement techniques of a given parameter from study to study, large coefficients of variation with repeated measurements or from study to study of a given parameter,^{111,112} different patient populations (eg, long-term vs short-term ventilator dependence),^{113,114} and the absence of objective criteria to determine a patient's tolerance for a trial of either discontinuation or extubation.

Third, assessed outcomes differ from study to study. Some investigators have examined successful tolerance of a spontaneous breathing trial (SBT), others have used permanent discontinuation of the ventilator, and others have combined successful discontinuation and extubation. This latter approach is not optimal, given the differences in the pathophysiology of discontinuation versus extubation failure (see below).^{102,106} In addition, different studies use different durations of ventilator discontinuation or extubation to define success or failure. Although 24–48 hours of unassisted breathing often is considered to define the successful discontinuation of ventilator support, many studies use shorter time periods to indicate success and often do not report subsequent reintubation rates or the need to reinstitute mechanical ventilatory support.

Fourth, a number of ways have been used to express predictor performance, and many can be confusing or misleading. Traditional indexes of diagnostic test power include sensitivity/specificity and positive/negative predictive values. These indexes are limited, however, in that they rely on a single cut point or threshold and that they do not provide an easy way to go from pretest likelihood or probability, through testing, to a posttest probability. The McMaster AHCPR report⁴ recommends the use of likelihood ratios (LRs), and these will be used in this report to describe predictor performance. The LR is an expression of the odds that a given test result will be present in a patient with a given condition compared to a patient without the condition. An LR > 1 indicates that the probability of success increases, while values < 1 indicate that the probability of failure increases. LRs between 0.5 and 2 indicate that a weaning parameter is associated with only small, clinically unimportant changes in the posttest probability of success or failure. In contrast LRs from 2 to 5 and from 0.3 to 0.5 correlate with small but potentially important changes in probability, while ratios of 5 to 10 or 0.1 to 0.3 correlate with more clinically important changes

Table 3. Criteria Used in Weaning/Discontinuation Studies to Determine Whether Patients Receiving High Levels of Ventilatory Support Can Be Considered for Discontinuation (ie, Entered Into the Trials)*

Objective Measurements	Adequate oxygenation (eg, $P_{O_2} \geq 60$ mm Hg on $F_{IO_2} \leq 0.4$; PEEP ≤ 5 –10 cm H_2O ; $P_{O_2}/F_{IO_2} \geq 150$ –300) Stable cardiovascular system (eg, HR ≤ 140 beats/min; stable blood pressure; no or minimal vasopressors) Afebrile (eg, temperature $< 38^\circ C$) No significant respiratory acidosis Adequate hemoglobin (eg, Hgb ≥ 8 –10 g/dL) Adequate mentation (eg, arousable, GCS ≥ 13 , no continuous sedative infusions) Stable metabolic status (eg, acceptable electrolytes)
Subjective Clinical Assessments	Resolution of disease acute phase; physician believes discontinuation possible; adequate cough

P_{O_2} = partial pressure of oxygen
 F_{IO_2} = fraction of inspired oxygen
PEEP = positive end-expiratory pressure
HR = heart rate
GCS = Glasgow Coma Scale

*Adapted from References 101–103, 107–109, 119, and 120

in probability. Ratios of > 10 or < 0.1 correlate with very large changes in probability.¹¹⁵

Finally, because some investigators report data as continuous values (eg, means) rather than providing defined threshold values, combining studies using meta-analytic techniques often cannot be done.

Recommendation 2. *Patients receiving mechanical ventilation for respiratory failure should undergo a formal assessment of discontinuation potential if the following criteria are satisfied:*

1. Evidence for some reversal of the underlying cause of respiratory failure;
2. Adequate oxygenation (eg, $P_{aO_2}/F_{IO_2} > 150$ –200; requiring positive end-expiratory pressure [PEEP] ≤ 5 –8 cm H_2O ; $F_{IO_2} \leq 0.4$ –0.5) and pH (eg, ≥ 7.25);
3. Hemodynamic stability as defined by the absence of active myocardial ischemia and the absence of clinically important hypotension (ie, a condition requiring no vasopressor therapy or therapy with only low-dose vasopressors such as dopamine or dobutamine $< 5 \mu g/kg/min$); and
4. The capability to initiate an inspiratory effort.

The decision to use these criteria must be individualized. Some patients not satisfying all of the above the criteria (eg, patients with chronic hypoxemia below the thresholds cited) may be ready for attempts at discontinuation of mechanical ventilation.

Rationale and Evidence (Grade B)

While some investigators argue that the process of discontinuation starts as soon as the patient is intubated, it would seem reasonable that an appropriate level of ventilatory support should be maintained until the underlying cause of acute respiratory failure and any complicating issues have shown some sign of reversal. Indeed, patients with unresolving respiratory failure who require high levels of ventilatory support are probably at high risk for respiratory muscle fatigue (and the consequent prolongation of the need for mechanical ventilation) if aggressive reductions in support are undertaken.^{50,52,116–118}

The criteria used by clinicians to define disease “reversal,” however, have been neither defined nor prospectively evaluated in a randomized controlled trial. Rather, various combinations of subjective assessment and objective criteria (eg, usually gas exchange improvement, mental status improvement, neuromuscular function assessments, and radiographic signs) that may serve as surrogate markers of recovery have been employed (Table 3).^{101–103,107–109,119,120} It should be noted, however, that some patients who have not ever met one or more of these criteria still have been shown to be capable of eventual liberation from the ventilator.¹⁰⁴

These “clinical assessments” of the status of the patient’s respiratory failure, however, are not enough to make decisions on the discontinuation of support. For example, one survey¹²¹ of intensivists using clinical judgment to assess the potential for discontinuation found a sensitivity of only 35% (6 of 17 patients who were successfully discontinued were identified) and a specificity of 79% (11 of 14 who failed discontinuation were identified). Moreover, in 2 large trials,^{107,109} despite the presence of apparent disease stability/reversal prior to performing a screening SBT, the managing clinicians did not recognize that discontinuation was feasible in almost two thirds of the subjects. Thus, the conclusion is that some evidence of “clinical” stability/reversal is a key first step in assessing for discontinuation potential but that more focused assessments are needed before deciding to continue or discontinue ventilatory support.

Recommendation 3. *Formal discontinuation assessments for patients receiving mechanical ventilation for respiratory failure should be done during spontaneous breathing rather than while the patient is still receiving substantial ventilatory support. An initial brief period of spontaneous breathing can be used to assess the capability of continuing onto a formal SBT. The criteria with which to assess patient tolerance during SBTs are the respiratory pattern, adequacy of gas exchange, hemodynamic stability, and subjective comfort. The tolerance of SBTs lasting 30 to*

EVIDENCE-BASED GUIDELINES FOR WEANING AND DISCONTINUING VENTILATORY SUPPORT

Table 4. Measurements Performed Either While Patient Was Receiving Ventilatory Support or During a Brief Period of Spontaneous Breathing That Have Been Shown to Have Statistically Significant LRs to Predict the Outcome of a Ventilator Discontinuation Effort in More Than One Study

Parameter	Studies (n)	Threshold Values	Positive LRs Range
<u>Measured on Ventilator</u>			
\dot{V}_E	20	10–15 L/min	0.81–2.37
NIF	10	–20 to –30 cm H ₂ O	0.23–2.45*
$P_{I_{max}}$	16	–15 to –30 cm H ₂ O	0.98–3.01
$P_{0.1}/P_{I_{max}}$	4	0.30	2.14–25.3
CROP score	2	13	1.05–19.74
<u>Measured During a Brief Period of Spontaneous Breathing</u>			
RR	24	30–38 breaths/min	1.00–3.89
V_T	18	325–408 mL (4–6 mL/kg)	0.71–3.83
f/ V_T ratio	20	60–105/L	0.84–4.67

LR = likelihood ratio
 \dot{V}_E = minute ventilation
 *1 study reported a likelihood ratio (LR) of 35.79.
 NIF = negative inspiratory force (maximum inspiratory pressure)
 $P_{I_{max}}$ = maximum inspiratory pressure
 $P_{0.1}$ = mouth occlusion pressure 0.1 s after the onset of inspiratory effort
 CROP = index including compliance, respiratory rate, oxygenation, and pressure
 RR = respiratory rate
 V_T = tidal volume
 f/ V_T = respiratory rate/tidal volume ratio

Table 5. Frequency of Tolerating an SBT in Selected Patients and Rate of Permanent Ventilator Discontinuation Following a Successful SBT

Study	Patients Receiving SBT	Patients Tolerating SBT n (%)	Patients Discontinuing Ventilation	Patients Having Ventilation Reinstated n (%)
Esteban et al ¹⁰⁵	546	416 (76)	372	58 (16)
Ely et al ¹⁰⁸	113	88 (78)	65	5 (4)
Dojat et al ¹¹⁰	38	22 (58)	22	5 (23)
Esteban et al ¹⁰¹	246	192 (78)	192	36 (19)
Esteban et al ¹⁰²	270*	237 (89)	237	32 (14)
Esteban et al ¹⁰²	256†	216 (84)	216	29 (13)

SBT = spontaneous breathing trial
 *30 min SBT
 †120 min SBT

120 minutes should prompt consideration for permanent ventilator discontinuation.

Rationale and Evidence (Grade A)

Because clinical impression is so inaccurate in determining whether or not a patient meeting the criteria listed in Table 3 will successfully discontinue ventilatory support, a more focused assessment of discontinuation potential is necessary. These assessments can be performed either during spontaneous breathing or while the patient is still receiving substantial ventilatory support. These assessments can be used not only to drive decisions on weaning and discontinuation (ie, functioning as predictors) but also to offer insight into mechanisms of discontinuation failures.

The McMaster AH CPR report⁴ found evidence in the literature supporting a possible role for 66 specific measurements as predictors. Some of these (eg, the negative effects of the duration of mechanical ventilation, and the length of/difficulty of surgery^{44,122–124}) were derived from general clinical observations, but most were from studies on focused assessments of the patient’s respiratory system. From these, the McMaster AH CPR group identified 8 parameters that had consistently significant LRs to predict successful discontinuation in several studies. Some of these measurements are made while the patient is still receiving ventilatory support; others require an assessment during a brief period of spontaneous breathing. These parameters, their threshold values, and the range of reported LRs are given in Table 4. It should be noted that despite the sta-

Table 6. Criteria Used in Several Large Trials* to Define Tolerance of an SBT

Objective measurements indicating tolerance/success	Gas exchange acceptability ($S_{pO_2} \geq 85\text{--}90\%$; $P_{O_2} \geq 50\text{--}60$ mm Hg; $pH \geq 7.32$; increase in $P_{aCO_2} \leq 10$ mm Hg)
	Hemodynamic stability (HR < 120–140 beats/min; HR not changed > 20%; systolic BP < 180–200 mm Hg and > 90 mm Hg; BP not changed > 20%, no pressors required)
	Stable ventilatory pattern (eg, RR $\leq 30\text{--}35$ breaths/min, RR not changed > 50%)
Subjective clinical assessments indicating intolerance/failure	Change in mental status (eg, somnolence, coma, agitation, anxiety)
	Onset or worsening of discomfort
	Diaphoresis
	Signs of increased work of breathing (use of accessory respiratory muscles, thoracoabdominal paradox)

SBT = spontaneous breathing trial

S_{pO_2} = oxygen saturation measured by pulse oximetry

P_{O_2} = partial pressure of oxygen

P_{aCO_2} = arterial partial pressure of carbon dioxide

HR = heart rate

RR = respiratory rate

*References 101, 102, 105, 108–110, 119, and 120.

tistical significance of these parameters, the generally low LRs indicate that the clinical applicability of these parameters alone to individual patients is low.

Although assessments that are performed while a patient is receiving substantial ventilatory support or during a brief period of spontaneous breathing (Tables 3 and 4) can yield important information about discontinuation potential, assessments that are performed during a formal, carefully monitored SBT appear to provide the most useful information to guide clinical decision-making regarding discontinuation. Indeed, because of the efficacy and safety of a properly monitored SBT (see below), the assessments in Table 4 that are performed to predict SBT outcome are generally unnecessary.

In concept, the SBT should be expected to perform well, as it is the most direct way to assess a patient's performance without ventilatory support. Indeed, the evidence for this concept is quite strong. As can be seen in Table 5, multiple studies have found that patients tolerant of SBTs that are 30 to 120 min in length were found to have successful discontinuations at least 77% of the time.

Because the 12 to 42% of patients in the Table 5 studies failing the SBT were not systematically removed from ventilatory support, the ability of a failed SBT to predict the need for ventilator dependence (ie, negative predictive

value) cannot be formally assessed. Indeed, it is conceivable that iatrogenic factors such as endotracheal tube discomfort or continuous positive airway pressure (CPAP) demand-valve insensitivity/unresponsiveness, rather than true ventilator dependence, caused the failure of the SBT in at least some of these patients.¹²⁵ Thus, it is thus unclear how many patients who are unable to tolerate an SBT would still be able to tolerate long-term ventilator discontinuation. Although the number is likely to be small, it is probably not zero, and this needs to be considered when dealing with patients who repeatedly fail an SBT.

The criteria used to define SBT "tolerance" are often integrated indexes since, as noted above, single parameters alone perform so poorly. These integrated indexes usually include several physiologic parameters as well as clinical judgment, incorporating such difficult-to-quantify factors as "anxiety," "discomfort," and "clinical appearance." The criteria that have been used in several large trials are given in Table 6.

A potential concern about the SBT is safety. Although unnecessary prolongation of a failing SBT conceivably could precipitate muscle fatigue, hemodynamic instability, discomfort, or worsened gas exchange,^{19,50,117,126,127} there are no data showing that SBTs contribute to any adverse outcomes if terminated promptly when failure is recognized. Indeed, in a cohort of > 1,000 patients in whom SBTs were routinely administered and properly monitored as part of a protocol, only one adverse event was thought to be even possibly associated with the SBT.¹⁰⁴

There is evidence that the detrimental effects of ventilatory muscle overload, if it is going to occur, often occurs early in the SBT.^{73,108,110,128} Thus, the initial few minutes of an SBT should be monitored closely, before a decision is made to continue (this is often referred to as the "screening" phase of an SBT). Thereafter, the patient should continue the trial for at least 30 min, but for not > 120 min,¹⁰² to assure maximal sensitivity and safety. It also appears that whether the SBT is performed with low levels of CPAP (eg, 5 cm H₂O), low levels of pressure support (eg, 5–7 cm H₂O), or simply as "T-piece" breathing has little effect on outcome.^{101,129–131} CPAP, however, conceivably could enhance breath triggering in patients with significant auto-PEEP.^{132,133}

Recommendation 4. *The removal of the artificial airway from a patient who has successfully been discontinued from ventilatory support should be based on assessments of airway patency and the ability of the patient to protect the airway.*

Rationale and Evidence (Grade C)

Extubation failure can occur for reasons distinct from those that cause discontinuation failure. Examples include upper airway obstruction or inability to protect the airway

and clear secretions. The risk of postextubation upper airway obstruction increases with the duration of mechanical ventilation, female gender, trauma, and repeated or traumatic intubation.¹⁰⁶ The detection of an air leak during mechanical ventilation when the endotracheal tube balloon is deflated can be used to assess the patency of the upper airway (cuff leak test).¹³⁴ In a study of medical patients,¹³⁵ a cuff leak < 110 mL (ie, average of 3 values on 6 consecutive breaths) measured during assist control ventilation within 24 hours of extubation identified patients at high risk for postextubation stridor. Although others have not confirmed the utility of the cuff leak test for predicting postextubation stridor,¹³⁶ many patients who develop this can be treated with steroids and/or epinephrine (and possibly with noninvasive ventilation and/or heliox) and do not necessarily need to be reintubated. Steroids and/or epinephrine could also be used 24 hours prior to extubation in patients with low cuff leak values. It is also important to note that a low value for cuff leak may actually be due to encrusted secretions around the tube rather than to a narrowed upper airway. Despite this, reintubation equipment (including tracheostomy equipment) should be readily available when extubating patients with a low cuff leak values.

The capacity to protect the airway and to expel secretions with an effective cough would seem to be vital for extubation success, although specific data supporting this concept are few. Successful extubations have been reported¹³⁷ in a select group of brain-injured comatose patients who were judged to be capable of protecting their airways. However, it is difficult to extrapolate this experience to more typical ICU patients, and many would argue that some capability of the patient to interact with the care team should be present before the removal of an artificial airway. Airway assessments generally include noting the quality of cough with airway suctioning, the absence of “excessive” secretions, or the frequency of airway suctioning (eg, every 2 h or more).^{34,108,138} Coplin et al¹³⁷ devised an “airway care score,” which semiquantitatively assesses cough, gag, suctioning frequency, and sputum quantity, viscosity, and character that predicted extubation outcome. Peak cough flows of > 160 L/min predict successful trans-laryngeal extubation or tracheostomy tube decannulation in neuromuscular- or spinal cord-injured patients.¹³⁹

Managing the Patient Who Has Failed a Spontaneous Breathing Test

Introduction

The failure of a patient to complete an SBT raises 2 important questions. First, what caused the SBT failure, and are there readily reversible factors that can be corrected? Second, how should subsequent mechanical ven-

tilatory support be managed? Specifically, should an SBT be tried again? If so, when? What form of ventilatory support should be provided in between SBTs and should support be at a constant high level or should efforts be made to routinely reduce the level of support gradually (ie, to wean support)?

Evaluating evidence addressing mechanical ventilatory support strategies is particularly problematic. This is because trials comparing 2 or more approaches to ventilator management compare not only the modes of ventilation but also how those modes are used. Ideally, trial design should be such that management philosophies and the aggressiveness of support reduction are similar in each strategy being evaluated. Unfortunately, this is often not the case, as investigator experience with one approach has a tendency to result in more favorable “rules” of support reduction for that approach compared to others.

Recommendation 5. *Patients receiving mechanical ventilation for respiratory failure who fail an SBT should have the cause for the failed SBT determined. Once reversible causes for failure are corrected, and if the patient still meets the criteria listed in Table 3, subsequent SBTs should be performed every 24 hours.*

Rationale and Evidence (Grade A)

Although failed SBTs are often a reflection of persistent respiratory system abnormalities,⁵² a failed SBT should prompt a search for other causes or complicating factors (see the “Pathophysiology of Ventilator Dependence” section). Specific issues include the adequacy of pain control, the appropriateness of sedation, fluid status, bronchodilator needs, the control of myocardial ischemia, and the presence of other disease processes that either can be readily addressed or else can be considered when deciding to proceed further with discontinuation attempts.

Assuming medical management is optimized and that the patient who has failed an SBT still meets the criteria listed in Table 3, the following 2 questions involving subsequent SBTs arise: First, should SBTs be attempted again or should another approach to ventilator withdrawal be attempted? Second, if an SBT is attempted again, when should that be?

There are some data on which to base an answer to the first question. The one large randomized trial¹⁰⁷ that compared routine SBTs to 2 other weaning strategies that did not include SBTs provides compelling evidence that SBTs administered at least once daily shorten the discontinuation period compared to strategies that do not include daily SBTs. In addition, 2 studies^{108,119} showing the success of protocol-driven ventilator discontinuation strategies over “usual care” both included daily SBTs. The subsequent use of routine subsequent SBTs in this patient population thus seems appropriate.

There are several lines of evidence that support waiting 24 hours before attempting an SBT again in these patients. First, except in patients recovering from anesthesia, muscle relaxants, and sedatives, respiratory system abnormalities rarely recover over a short period of hours and thus frequent SBTs over a day may not be expected to be helpful. Supporting this are data from Jubran and Tobin⁵² showing that failed SBTs often are due to persistent respiratory system mechanical abnormalities that are unlikely to reverse rapidly. Second, there are data suggesting that a failed SBT may result in some degree of respiratory muscle fatigue.^{50,117,118} If so, studies^{126,140} conducted in healthy subjects suggest that recovery may not be complete for anywhere from several hours to > 24 hours. Third, the trial by Esteban et al¹⁰⁷ specifically addressed this issue and provided strong evidence that twice-daily SBTs offer no advantage over a single SBT and, thus, would serve only to consume unnecessary clinical resources.

Recommendation 6. *Patients receiving mechanical ventilation for respiratory failure who fail an SBT should receive a stable, nonfatiguing, comfortable form of ventilatory support.*

Rationale and Evidence (Grade B)

There are a number of ventilator modes that can provide substantial ventilatory support as well as the means to reduce partial ventilatory support in patients who have failed an SBT (Table 7). A key question, however, is whether attempts at gradually lowering the level of support (weaning) offer advantages over a more stable, unchanging level of support between SBTs. The arguments for using gradual reductions are (1) that muscle conditioning might occur if ventilatory loads are placed on the patient's muscles and (2) that the transition to extubation or to an SBT might be easier from a low level of support than from a high level of support. Data supporting either of these claims, however, are few. However, maintaining a stable level of support between SBTs reduces the risk of precipitating ventilatory muscle overload from overly aggressive support reduction. It also offers a substantial resource consumption advantage in that it requires far less practitioner time. The study by Esteban et al¹⁰⁷ partially addressed this issue in that it compared daily SBTs (and a stable level of support in those who failed) to 2 other approaches using gradual reductions in support (ie, weaning with pressure support or intermittent mandatory ventilation [IMV]) and demonstrated that the daily SBT with stable support between tests permitted the most rapid discontinuation. What has not been addressed, however, is whether gradual support reductions *coupled* with daily SBTs offer any advantages.

The McMaster AHCPR report⁴ identified 3 other randomized trials^{109,141,142} that compared gradual reduction

Table 7. Modes of Partial Ventilator Support

Mode	Patient Work Adjusted By
SIMV	Number of machine breaths supplied (ie, the fewer the number of machine breaths, the more spontaneous breaths are required)
PSV	Level of inspiratory pressure assistance with spontaneous efforts
SIMV + PSV	Combining the adjustments of SIMV and PSV
VS	PSV with a "guaranteed" minimum V_T (PSV level adjusts automatically according to clinician V_T setting)
VAPS(PA)	PSV with "guaranteed" minimum V_T (additional flow is supplied at end inspiration if necessary to provide clinician V_T setting)
MMV	SIMV with a "guaranteed" \dot{V}_E (machine breath rate automatically adjusts according to clinician \dot{V}_E setting)
APRV	Pressure difference between inflation and release (ie, the less the pressure difference, the more spontaneous breaths are required)

SIMV = synchronized intermittent mandatory ventilation

PSV = pressure support ventilation

VS = volume support

VAPS(PA) = volume-assured pressure support (pressure augmentation)

V_T = tidal volume

MMV = mandatory minute ventilation

\dot{V}_E = minute ventilation

APRV = airway pressure-release ventilation

strategies using different modes but *not* routine daily SBTs. The study by Brochard et al¹⁰⁹ was the most similar in design to the study by Esteban et al¹⁰⁷ that was noted before, and it included a pressure-support group and an IMV group. A third group received gradually increasing fixed periods of spontaneous breathing that were designed only to provide brief periods of work and not specifically to test for discontinuation (ie, they were not routine daily SBTs, as defined above). The results showed that their use of gradually lengthening spontaneous breathing periods was inferior to other strategies and, like the Esteban trial, the pressure support strategy was easier to reduce than the IMV strategy. The other 2 randomized trials^{141,142} that were identified by the McMaster AHCPR report were much smaller than the Esteban et al¹⁰⁷ and Brochard et al,¹⁰⁹ and both suggested that pressure support was easier to reduce than IMV alone. Because none of these studies offer evidence that gradual support strategies are superior to stable support strategies between SBTs, the clinical focus for the 24 hours after a failed SBT should be on maintaining adequate muscle unloading, optimizing comfort (and thus sedation needs), and avoiding complications, rather than aggressive ventilatory support reduction.

Ventilator modes and settings can affect these goals.¹⁴³ Assisted modes of ventilation (as opposed to machine-controlled modes) are generally preferable in this setting

because they allow patient muscle activity and some patient control over the ventilatory pattern. Although good clinical supporting data are lacking, these features may help to avoid muscle disuse atrophy²⁴ and may reduce sedation needs in these types of patients.¹⁴³ With assisted modes, sensitive/responsive ventilator triggering systems,^{144–147} applied PEEP in the presence of a triggering threshold load from auto-PEEP,^{132,133} flow patterns matched to patient demand,^{148–152} and appropriate ventilator cycling to avoid air trapping^{153,154} are all important to consider in achieving patient comfort and minimizing imposed loads.

In recent years, several ventilator support modes (volume support,¹⁵⁵ adaptive support ventilation (ASV),^{155,156} minimum minute ventilation (MMV),¹⁵⁵ and a knowledge-based system for adjusting pressure support^{110,157}) have been developed in an attempt to “automatically” wean patients by feedback from one or more ventilator-measured parameters. Minimum minute ventilation set at either 75% of measured minute ventilation¹⁵⁸ or to a carbon dioxide target,¹²⁰ and a knowledge-based system for adjusting pressure support^{110,157} all have been shown to be capable of automatically reducing support safely in selected populations. However, none of these has been compared to the daily SBT approach described above. Moreover, the premises underlying some of these feedback features (eg, that an ideal volume can be set for volume support or that an ideal ventilatory pattern based on respiratory system mechanics can be set for ASV) may be flawed, especially in sick patients. Indeed, potentially flawed feedback logic may, in fact, delay support reduction. Further work is clearly needed to establish the role (if any) of these automated approaches.

There has been increasing interest in the use of noninvasive positive-pressure ventilation (NPPV) in recent years. Although NPPV has been used primarily as a method to avoid intubation, it has also been used as a technique to facilitate the discontinuation of invasive ventilatory support. Data from the pooling of results of 2 prospective, randomized controlled trials^{159,160} in patients with chronic respiratory disease suggest the need for reductions in the durations of mechanical ventilation, ICU stay, mortality, and the incidence of nosocomial pneumonia with postextubation support provided by NPPV. Appropriate patient selection and the feasibility of the widespread application of these findings remains to be determined.

Recommendation 7. *Anesthesia/sedation strategies and ventilator management aimed at early extubation should be used in postsurgical patients.*

Rationale and Evidence (Grade A)

The postsurgical patient poses unique problems for ventilator discontinuation. In these patients, depressed respi-

ratory drive and pain issues are the major reasons for ventilator dependence. Optimal sedation, pain management, and ventilator strategies offer opportunities to shorten the duration of mechanical ventilation.

The McMaster AHCPR report⁴ identified 5 randomized controlled trials in postcardiac surgery patients^{161–165} that demonstrated that a lower anesthetic/sedation regimen permitted earlier extubation. The pooled results showed a mean effect of 7 hours. Similar effects were found using these approaches in other postsurgical populations.^{166–170}

Ventilator modes that guarantee a certain breath rate and minute ventilation (assist control modes, IMV, and MMV) are important in patients with unreliable respiratory drives. However, frequent assessments and support reductions are necessary since recovery in these patients usually occurs over only a few hours. Aggressive support reduction strategies have been shown to lead to earlier discontinuations of ventilation.^{166,169} Conceptually, the immediate postoperative patient might be ideally suited for simple automatic feedback modes that provide a backup form of support (eg, MMV or ASV).^{120,156,158} Data showing improved outcomes or lower costs with these automated approaches, however, are lacking.

Recommendation 8. *Weaning/discontinuation protocols designed for nonphysician health care professionals (HCPs) should be developed and implemented by ICUs. Protocols aimed at optimizing sedation should also be developed and implemented.*

Rationale and Evidence (Grade A)

There is clear evidence that nonphysician HCPs (eg, respiratory therapists and nurses) can execute protocols that enhance clinical outcomes and reduce costs for critically ill patients.¹⁷¹ In recent years, 3 randomized controlled trials incorporating 1,042 patients also have demonstrated that outcomes for mechanically ventilated patients who were managed using HCP-driven protocols were improved over those of control patients managed with standard care. Specifically, Ely et al¹⁰⁸ published the results of a 2-step protocol driven by HCPs using a daily screening procedure followed by an SBT in those who met screening criteria. The discontinuation of mechanical ventilation was then recommended for patients tolerating the SBT. Although the 151 patients managed with the protocol had a higher severity of illness than the 149 control subjects, they were removed from the ventilator 1.5 days earlier (with 2 days less weaning), had 50% fewer complications related to the ventilator, and had mean ICU costs of care that were lower by > \$5,000 per patient. In a slightly larger trial with a more diverse patient population, Kollef et al¹¹⁹ used 3 different nonphysician-HCP-driven protocols and showed that the mean duration of mechanical

ventilation could be reduced by 30 hours. Finally, Marelich et al¹⁷² showed that the duration of mechanical ventilatory support could be reduced almost 50% using nurse-driven and therapist-driven protocols ($p = 0.0001$).

The reproducibility of benefit for using various protocols in different ICUs and institutions suggests that it is the use of a standardized approach to management rather than any specific modality of ventilator support that improves outcomes. Indeed, when other key features in the management of mechanically ventilated patients, such as sedation and analgesia, also are subjected to protocols, further reductions in the time spent receiving mechanical ventilation can be achieved. For example, in a randomized controlled trial¹⁷³ of a nursing-implemented sedation protocol for 321 patients receiving mechanical ventilation, the use of the protocol was associated with a 50% reduction in the duration of mechanical ventilation and 2-day and 3-day reductions in the median ICU and lengths of hospital stay, respectively (all p values < 0.01). More recently, Kress et al¹⁷⁴ published the results of a randomized controlled trial of 128 patients showing that a daily spontaneous awakening trial was associated with 2 days fewer spent receiving mechanical ventilation ($p = 0.004$) and a 3-day shorter ICU stay ($p = 0.02$).

The data do not support endorsing any one ventilator discontinuation protocol, and the choice of a specific protocol is best left to the individual institution. In designing these protocols, consideration should be given to other recommendations in this document as well as to the specific patient populations. For instance, medical patients with severe lung injury might benefit from one type of management strategy (see recommendations 2 to 5), whereas surgical patients recovering from anesthesia might benefit from another strategy (see recommendation 7). In the context of emerging data about the benefits of NPPV^{159,160} and the substantial roles of HCPs in providing this treatment, there should be efforts made to develop HCP-driven protocols for this modality.

While each institution must customize the protocols to local practice, there are important general concepts that may ease the process of implementation and enhance success.

First, protocols should not be used to replace clinical judgment, but rather to complement it. Protocols are meant as guides and can serve as the general default management decision unless the managing clinician can justify a departure from the protocol. Any such departure should be carefully assessed and used to guide possible future modifications of the protocol.

Second, protocols should not be viewed as static constructs, but rather as dynamic tools in evolution, which can be modified to accommodate new data and/or clinical practice patterns. More studies regarding the impact of protocol-based ventilator management are needed for specific

patient populations (eg, neurosurgical patients¹⁷⁵ and trauma patients¹⁷⁶), in specific organizational structures (eg, open vs closed units and teaching vs community hospitals), and using computer-assisted decision-making.

Third, institutions must be prepared to commit the necessary resources to develop and implement protocols.¹⁷⁷ For instance, the effective implementation of protocols requires adequate staffing, as it has been shown that if staffing is reduced below certain thresholds, clinical outcomes may be jeopardized.^{178,179} Indeed, in the specific context of the discontinuation of mechanical ventilation, reductions in nurse/patient ratios have been associated with prolonged duration of mechanical ventilation.¹⁸⁰

Role of Tracheotomy in Ventilator-Dependent Patients

Introduction

Tracheotomy is commonly performed for critically ill, ventilator-dependent patients to provide long-term airway access. The benefits commonly ascribed to tracheotomy, compared to prolonged translaryngeal intubation, include improved patient comfort, more effective airway suctioning, decreased airway resistance, enhanced patient mobility, increased opportunities for articulated speech, ability to eat orally, and a more secure airway. Conceptually, these advantages might result in fewer ventilator complications (eg, ventilator-associated pneumonia), accelerated weaning from mechanical ventilation, and the ability to transfer ventilator-dependent patients from the ICU. Concern, however, exists about the risks associated with the procedure and the costs involved.

The impact of tracheotomy on the duration of mechanical ventilation and ICU outcomes in general has been examined by several different study designs—none of them ideal. Most studies are retrospective, although a few prospective studies have been performed. A serious problem is that many studies assigned patients to treatment groups on the basis of physician practice patterns rather than random assignment. Those studies that used random assignment frequently used quasi-randomization methods (eg, every other patient, every other day, hospital record number, odd-even days). Studies have compared patients undergoing tracheotomy versus those not undergoing tracheotomy and patients undergoing early tracheotomy versus those undergoing late tracheotomy. The definition of early versus late tracheotomy varies between studies. “Early” may be defined as a period as short as 2 days to as late as 10 days after the start.

Patient populations included in studies also vary widely between investigations and include general surgical and medical patients in some studies and specific patient groups

(eg, trauma patients or head-injured patients) in other studies. Most studies have design flaws in the collection and analysis of data, foremost of which is the absence of blinding. The absence of blinding is especially important considering that no study has used explicit, systematic protocols for weaning to control for any effects of tracheotomy on altering the approaches of clinicians to weaning. Finally, an outcome such as transfer to a non-ICU setting may depend on local resources, such as the availability of a non-ICU ventilator service.

Because there is such a surprisingly small amount of quality data regarding the relative impact of tracheotomy in terms of patient outcome relative to prolonged translaryngeal intubation, past recommendations for timing the procedure to achieve these benefits have been based on expert consensus.¹⁸¹

Recommendation 9. *Tracheotomy should be considered after an initial period of stabilization on the ventilator when it becomes apparent that the patient will require prolonged ventilator assistance. Tracheotomy should then be performed when the patient appears likely to gain one or more of the benefits ascribed to the procedure. Patients who may derive particular benefit from early tracheotomy are the following:*

- *Those requiring high levels of sedation to tolerate translaryngeal tubes*
- *Those with marginal respiratory mechanics (often manifested as tachypnea) in whom a tracheostomy tube having lower resistance might reduce the risk of muscle overload*
- *Those who may derive psychological benefit from the ability to eat orally, communicate by articulated speech, and experience enhanced mobility; and*
- *Those in whom enhanced mobility may assist physical therapy efforts*

Rationale and Evidence (Grade B)

While carrying some risks, tracheotomies in ventilator-dependent patients are generally safe. The problems associated with tracheotomy include perioperative complications related to the surgery, long-term airway injury, and the cost of the procedure.

Patient series reported during the early 1980s¹⁸² suggested that tracheotomy had a high risk of perioperative and long-term airway complications such as tracheal stenosis. More recent studies,^{183–185} however, have established that standard surgical tracheotomy can be performed with an acceptably low risk of perioperative complications. Regarding long-term risks, analyses of longitudinal studies suggest that the risk of tracheal stenosis after tracheotomy is not clearly higher than the

risks of subglottic stenosis from prolonged translaryngeal intubation.¹⁸⁶ Also, the nonrandomized studies commonly reported in the literature bias results toward greater long-term airway injury in patients undergoing tracheotomy because the procedure was performed after a prolonged period of translaryngeal intubation, which may prime the airway for damage from a subsequent tracheotomy.^{182,187–189} Finally, the cost of tracheotomy can be lowered if it is performed in the ICU rather than an operating room, either by the standard surgical or percutaneous dilational technique.^{186,187} Even when tracheotomy is performed in an operating room, the cost may be balanced by cost savings if a ventilator-dependent patient can be moved from an ICU setting after the placement of a tracheostomy. The actual cost benefits of tracheotomy, however, have not been established because no rigorous cost-effectiveness analyses have been performed.

Given the above conditions, it seems reasonable to conclude that none of the potential problems with tracheotomy is of sufficient magnitude to make tracheotomy any less clinically acceptable compared with other procedures commonly performed in critically ill patients.

Potentially, the most important beneficial outcome from a tracheotomy would be to facilitate the discontinuation of mechanical ventilatory support. Supporting evidence comes both from observations on “intermediate” end points (eg, comfort and mobility, decreased airway resistance, and a lower incidence of ventilator-associated pneumonia) as well as ICU outcome studies examining the duration of mechanical ventilation, ICU length of stay (LOS), and mortality. This evidence is reviewed below.

Improved Patient Comfort: No prospective outcome studies in general populations of ventilator-dependent patients using validated measurement tools have established that tracheotomy results in greater patient comfort or mobility, compared with prolonged translaryngeal intubation. Indeed, to our knowledge, only one study¹⁸³ has attempted to document this by reporting that interviewed ICU caregivers believed ventilated patients were more comfortable after tracheotomy. Despite this lack of data, the general clinical consensus is that patients supported with long-term mechanical ventilation have less facial discomfort when nasotracheal or orotracheal endotracheal tubes are removed and a tracheotomy is performed. Furthermore, patient well-being is thought to be promoted by a tracheotomy through its effects on assisting articulated speech, oral nutrition, and mobility, which may promote the discontinuation of sedatives and analgesics. The maintenance of continuous sedation has been associated with the prolongation of mechanical ventilation,¹⁹⁰ but the effects of tracheotomy on sedation usage have not been studied specifically.

Table 8. The Impact of Tracheotomy on Mechanical Ventilation Outcome

Study	Patient Type (n)	Design	Outcome
Lesnik et al ²⁰¹	Trauma/101	Randomized early (6 d) vs late (21 d) tracheotomy	Vent days shorter with earlier tracheotomy
Blot et al ²⁰²	Neutropenic/53	Retrospective early (2 d) vs late (> 7 d) tracheotomy or no tracheotomy	LOS longer in patients undergoing early tracheotomy
Koh et al ²⁰³	Neurosurgical/49	Retrospective elective (9 d) vs failed extubation tracheotomy	LOS shorter in elective tracheotomy
Dunham and LaMonica ¹⁹⁹	Trauma/74	Prospective early (4 d) vs late (14 d) or no tracheotomy	No effect
El Nagger et al ²⁰⁴	General acute respiratory failure/52	Prospective (3 d) vs delayed (> 10 d) tracheotomy	More patients weaned in delayed tracheotomy
Rodriguez et al ¹⁹⁸	Trauma/106	Randomized early (\leq 7 d) vs late (> 7 d) tracheotomy	LOS shorter in patients undergoing early tracheotomy but those weaned before late tracheotomy not considered
Sugerman et al ²⁰⁰	Trauma/126	Randomized early (3–5 d) vs late (10–21 d) tracheotomy	No effect

LOS = length of stay

Decreasing Airway Resistance: Although the small radius of curvature of tracheostomy tubes increases turbulent air flow and airway resistance, the short length of tracheostomy tubes results in an overall lowering of airway resistance (and thus reduced patient muscle loading) when compared to standard endotracheal tubes in both laboratory and clinical settings.^{191–197} While the development of secretions will increase resistance in both tracheostomy and endotracheal tubes, easier suctioning and removable inner cannulas may reduce this effect in tracheostomy tubes.¹⁹⁵ The existing data thus indicate that airway resistance and muscle loading may decrease in some patients after the performance of tracheotomy, but the clinical impact of this improvement in pulmonary mechanics on weaning has not been established. Conceivably, patients with borderline pulmonary mechanics may benefit from tracheotomy because of decreased airway resistance, which becomes more clinically important with high respiratory rates.

Impact of Tracheotomy on Ventilator-Associated Pneumonia: Early tracheotomy and, alternatively, the avoidance of tracheotomy by maintaining a translaryngeal endotracheal tube in place both have been proposed as strategies to promote the successful discontinuation of mechanical ventilation by avoiding ventilator-associated pneumonia. Few data support the conclusion, however, that the timing of tracheotomy alters the risk of pneumonia. Three prospective studies have evaluated the relative risk of pneumonia in patients randomized to early versus late tracheotomy.^{198–200} These studies examined 289 patients and found a relative risk for pneumonia (early tracheotomy vs late tracheotomy group) of only 0.88 (95% confidence

interval 0.70 to 1.10). Considerable methodological flaws in these studies, however, do not allow firm conclusions to be drawn regarding the effects of tracheotomy on pneumonia risk. Presently, no data support the competing contentions that early tracheotomy either decreases or increases the risk of ventilator-associated pneumonia.

Outcome Studies: Impact of Tracheotomy on Duration of Mechanical Ventilation: The results of a number of studies examining ICU outcome (ie, ventilator days, ICU LOS, mortality) have been reported, and are summarized in Table 8.^{198–204} Several of these studies were appraised in a systematic review.²⁰⁵ The authors of this review concluded that insufficient evidence existed to support the contention that the timing of tracheotomy alters the duration of mechanical ventilation in critically ill patients. Also, the review identified multiple flaws in the available studies. There appears to be a clinical impression that timing of tracheotomy promotes discontinuation of mechanical ventilation in some ventilator-dependent patients, but not all. However, the quality of existing studies does little to support this clinical impression.

In the future, because of the difficulty in blinding caregivers to the presence or absence of tracheotomy, studies should use explicit weaning protocols to control for different levels of approaches toward weaning that the presence of a tracheotomy may invoke. Studies also could be improved by more rigorous patient inclusion and exclusion criteria, better accounting for dropouts, the use of conventional randomization methods, multi-center designs to allow sufficient sample sizes to determine the interaction of underlying conditions, and multivariate analysis techniques. Cost-effectiveness analysis

Table 9. Comparison of Observational Studies*

Variable	Indihar ²¹³	Latriano et al ²¹⁴	Petrak et al ²¹⁵	Clark and Theiss ²¹⁶	Bagley and Cooney ²¹⁷	Scheinhorn et al ²⁰⁸	Carson et al ²¹⁸	Dasgupta et al ²¹⁹	Gracey et al ²²⁰
Type of unit	NRCU	NRCU	RWC	RWC	RWC	RWC	RWC	NRCU	NRCU
Patients, No.	171	224	388	113	278	1,123	133	212	420
Age (y) [†]	–	67	72 [‡]	65	67	69	71	68	67 [§]
Sex (% women)	54	47	53	56	53	57	52	55	52 [§]
Diagnoses precipitating ventilator dependency	COPD	medical > surgical	medical > surgical	medical = surgical	medical > surgical	medical > surgical	medical > surgical	medical = surgical	surgical > medical
ICU days [†] receiving ventilation	55	23	42	–	–	44	25	25 [‡]	37 [§]
Days [†] to weaning in ICU	39	43	–	–	43	39	–	13 [‡]	10
% Weaned	34	51	51	47	38	56	38	60	60
% Survival to discharge	60	50	66	61	53	71	50	82	94
% Survival 12 mo after discontinuing ventilation	–	–	–	–	–	38	23	–	53 [¶]

*Each with > 100 patients transferred for weaning from prolonged mechanical ventilation (> 21 prior ICU ventilator days).

NRCU = noninvasive respiratory care unit

RWC = regional weaning center

[†]Mean

[‡]Median

[§]Data from an earlier report on the same cohort of patients.²¹¹

^{||}Authors' data were recalculated so that "% weaned" signifies number weaned/number of patients who were admitted to the hospital ventilator-dependent

[¶]Value attained 4 years after discharge

Adapted from Reference 224.

also would assist the determination of the value of tracheotomy for weaning.

The Role of Long-Term Facilities

Introduction

The patient who remains ventilator-dependent despite maximal medical/surgical therapy and aggressive attempts to remove ventilator support is becoming an increasing challenge for critical care practitioners. In 2 studies,^{206,207} up to 20% of medical ICU patients met the 21-day Health Care Financing Administration definition of prolonged mechanical ventilation (PMV).²⁰⁸ Better treatment and technology, no doubt, are playing a major role in the conversion of patients who would have died a decade ago into patients who survive today, but who have substantial remaining respiratory dysfunction.

Prior to the 1980s, these patients simply remained in ICUs and were managed using acute-care principles. The only other option was permanent ventilatory support in either the patient's home or in a nursing home. Financial pressures, coupled with the concept that the aggressive ICU mindset might not be optimal for the more slowly recovering patient, have led to creation of weaning facilities (both free-standing facilities and units within hospitals) that are potentially more cost-effective and better suited to meet the needs of these patients. A body of literature now is emerging that suggests that many patients

who previously would have been deemed "unweanable" may achieve ventilator independence in such facilities.

Recommendation 10. *Unless there is evidence for clearly irreversible disease (eg, high spinal cord injury or advanced amyotrophic lateral sclerosis), a patient requiring prolonged mechanical ventilatory support for respiratory failure should not be considered permanently ventilator-dependent until 3 months of weaning attempts have failed.*

Rationale and Evidence (Grade B)

A critical clinical issue is determining whether a patient requiring PMV has any hope of ventilator discontinuation or whether he/she is to have lifelong ventilator dependence. Patients in the former category clearly need attempts at ventilator discontinuation to be pursued, whereas patients in the latter category are only being subjected to unnecessary episodes of worsening respiratory failure with such attempts. These latter patients, instead, need to have the clinical focus changed to establish a lifelong support program.

Data from a number of centers caring for the patient requiring long-term ventilation offer insight into this question. In the Barlow Regional Weaning Center experience,^{208–210} patients with prolonged ventilator dependence following acute cardiorespiratory failure were still undergoing ventilator discontinuation up to 3 months (and, on occasion, 6 months) postintubation. Other studies^{211,212} suggest similar results in postsurgical and medical populations. Data from these stud-

Table 10. Post-ICU Weaning Strategies

Study	Strategies to Decrease Ventilator Support to Approximately Half Full Support	Strategies to Increase SBT
Gracey et al ^{220*}	IMV/PSV	TTO
Scheinhorn et al ²⁰⁸	IMV/PSV	VTM
Bagley and Cooney ²¹⁷	–	VTM/T-piece
Petrak et al ^{215*}	AC or PSV	VTM
Clark and Theiss ²¹⁶	PSV	VTM/T-piece
Latriano et al ²¹⁴	IMV/PSV	VTM/T-piece

ICU = intensive care unit
 SBT = spontaneous breathing trial
 *Data from personal communication
 IMV = intermittent mandatory ventilation
 PSV = pressure-support ventilation
 TTO = transtracheal oxygen
 VTM = Venti-Trach mask (tracheal collar)
 AC = assist-control

ies on the time that patients spend, on average, dependent on ventilator support in the ICU (36 d) and during subsequent weaning in the post-ICU setting (31 d), suggest a time frame for the reasonable continuance of ventilator discontinuation attempts. Thus the weight of evidence is that several months of attempts at ventilator discontinuation are required before most patients who are receiving mechanical ventilation for acute respiratory failure can be declared to be permanently ventilator-dependent.

Recommendation 11. *Critical-care practitioners should familiarize themselves with facilities in their communities, or units in hospitals they staff, that specialize in managing patients who require prolonged dependence on mechanical ventilation. Such familiarization should include reviewing published peer-reviewed data from those units, if available. When medically stable for transfer, patients who have failed ventilator discontinuation attempts in the ICU should be transferred to those facilities that have demonstrated success and safety in accomplishing ventilator discontinuation.*

Rationale and Evidence (Grade C)

There are >30 studies on post-ICU weaning from PMV. Most of these studies are observational, a reflection of the inherent logistical difficulties in designing randomized controlled trials in such a heterogeneous population that is being treated in such diverse settings. Table 9 summarizes the studies that report outcomes in > 100 patients in which PMV is defined as > 21 days of ventilator dependency. Most of these studies^{208,211–219} support the conclusion that ICU patients receiving PMV can be discontinued effectively and safely from ventilation when they are transferred to units dedicated to that activity. Efficacy is suggested in the fact that the data in Table 9 encompass 3,062

post-ICU patients and show that 1,588 (52%) successfully discontinued mechanical ventilation. Safety is suggested in that a 69% overall survival-to-discharge rate in this “chronically critically ill” cohort seems quite reasonable.

The facilities generating the data of Table 9 are of 2 basic types: (1) Most facilities, but not all, are licensed as long-term acute-care hospitals, which are required by the Health Care Financing Administration to maintain a mean LOS of > 25 days. These facilities are most often free-standing hospitals, which may have their own ICUs. Called “regional weaning centers” in Table 9, they serve several to many hospitals in their geographic areas. (2) Step-down units or noninvasive respiratory care units have no specific LOS requirement. These units usually reside within a host hospital and primarily serve that hospital. While both settings have acute-care staffing, but not critical care (ICU) staffing, they are often dissimilar in hospital admission and discharge criteria, treatment capability, and the availability of specialty/subspecialty consultation services and procedures offered on site, all of which may have a significant effect on outcome of care. Both of these types of facilities are characterized by less intensive staffing and less costly monitoring equipment, and, therefore, they generate less cost per patient than do ICUs.^{219,220}

Recommendation 12. *Weaning strategy in the PMV patient should be slow-paced and should include gradually lengthening self-breathing trials.*

Rationale and Evidence (Grade C)

Despite differences in patient population and physical facilities, the available information on strategies for discontinuation of PMV in the studies listed in Table 9 has a number of similarities (Table 10). Daily SBTs initially are uncommon since patients have already established themselves as very unlikely to “turn around” in 24 hours. Instead, ventilator support is gradually reduced, using common modes of partial support (Table 7). Usually at the point of approximately half-support, patients are switched to the SBT approach described above, which often is employed along with self-breathing trials of increasing duration. Since most patients are tracheotomized, tracheal collars are used, instead of the familiar T-piece in the ICU, to supply oxygen and humidity. During these procedures, it is important for the staff to remain patient. Psychological support and careful avoidance of unnecessary muscle overload is important for these types of patients.

Long-term facilities may be a particularly useful place to implement protocols such as described earlier in the “Managing the Patient Who Has Failed an SBT” section (Recommendation 8).^{221–223} Skilled nonphysician personnel (eg, registered nurses and registered respiratory therapists) are generally present in these units continually and, thus, are in a position to constantly interact with the patient

and to make ventilator adjustments as appropriate. Of interest, after the implementation of a therapist-implemented protocol for weaning, the time to wean in a regional weaning center, with its predominantly "medical" population, declined from a median of 29 days to < 17 days over a 2-year period of protocol use.^{22,3}

REFERENCES

- Manthous CA. Liberation from mechanical ventilation, Parts I & II. *Respir Care Clin North Am* 2000;6(2 & 3).
- Tobin MJ, Alex CG. Discontinuation of mechanical ventilation. In: Tobin MJ, editor. *Principles and practice of mechanical ventilation*. New York: McGraw-Hill 1994;1177-1206.
- Esteban A, Alia I, Ibanez J, Benito S, Tobin MJH. Modes of mechanical ventilation and weaning: a national survey of Spanish hospitals. The Spanish Lung Failure Collaborative Group. *Chest* 1994;106:1188-1193.
- Cook D, Meade M, Guyatt G, Griffith L, Booker L. Evidence report on criteria for weaning from mechanical ventilation. *AHCPR*: Nov, 1999.
- Raju P, Manthous CA. The pathogenesis of respiratory failure. *Respir Clin N Am* 2000;6(2):195-212.
- Fishman AP, Turino GM, Bergofsky EH. The syndrome of alveolar hypoventilation. *Am J Med* 1957;23:333-339.
- Devereaux MW, Keane JR, Davis RL. Automatic respiratory failure associated with infarction of the medulla. *Arch Neurol* 1973; 29:46-52.
- Hughes JMB. Central respiratory failure reversed by treatment. *Brain* 1967;90:675-680.
- Vallverdu I, Mancebo J. Approach to patients who fail initial weaning trials. *Respir Care Clin N Am* 2000;6(3):365-384.
- Lopata M, Onal E. Mass loading, sleep apnea, and the pathogenesis of obesity hypoventilation. *Am Rev Respir Dis* 1982;126:640-645.
- Onal E, Lopata M. Periodic breathing and pathogenesis of occlusive sleep apneas. *Am Rev Respir Dis* 1982;128:676-680.
- Daum JJ, Canter HG, Katz S. Central nervous system sarcoidosis with alveolar hypoventilation. *Am J Med* 1965;38:893-898.
- Barrientos-Vega R, Mar Sanchez-Soria M, Morales-Garcia C, et al. Prolonged sedation of critically ill patients with midazolam or propofol: impact on weaning and costs. *Crit Care Med* 1997;25:33-40.
- Wheeler AP. Sedation, analgesia, and paralysis in the intensive care unit. *Chest* 1993;104(2):566-577.
- Diehl JL, Lofaso F, Deleuze P, et al. Clinical relevant diaphragmatic dysfunction after cardiac operations. *J Thoracic Cardiovasc Surg* 1994;107:487-498.
- Hansen-Flaschen JH, Cowen J, Raps EC, et al. Neuromuscular blockade in the intensive care unit: more than we bargained for. *Am Rev Respir Dis* 1993;147:234-236.
- Segredo V, Caldwell JE, Matthey MA, et al. Persistent paralysis in critically ill patients after long administration of vecuronium. *New Eng J Med* 1992;327:524-528.
- Bellemare F, Grassino A. Effect of pressure and timing of contraction on human diaphragm fatigue. *J Appl Physiol* 1982;53:1190-1195.
- Jubran A, Tobin MJ. Passive mechanics of lung and chest wall in patients who failed or succeeded in trials of weaning. *Am J Respir Crit Care Med* 1997;155:916-921.
- Roussos C, Macklem PT. The respiratory muscles. *N Engl J Med* 1982;307:786-797.
- Tobin MJ, Laghi F, Jubran A. Respiratory muscle dysfunction in mechanically ventilated patients. *Mol Cell Biochem* 1998;179(1-2):87-98.
- NHLBI Workshop summary. Respiratory muscle fatigue: report of the respiratory muscle fatigue workshop group. *Am Rev Respir Dis* 1990;142:474-480.
- Tobin MJ, Laghi F. Monitoring respiratory muscle function. In: Tobin MJ, editor. *Principles and practice of intensive care monitoring*. New York: McGraw-Hill 1998:497-544.
- Le Bourdelles G, Viïres N, Boezkowski J, et al. Effects of mechanical ventilation on diaphragmatic contractile properties in rats. *Am J Respir Crit Care Med* 1994;149:1539-1544.
- Bolton CF, Breuer AC. Critical illness polyneuropathy. *Muscle Nerve* 1999;22(3):419-424.
- Hubmayr RD, Rehder K. Respiratory muscle failure in critically ill patients. *Semin Respir Med* 1992;13:14-21.
- Reid WD, Huang J, Bryson S, Walker DC, Belcastro AN. Diaphragm injury and myofibrillar structure induced by resistive loading. *J Appl Physiol* 1994;76:176-184.
- Rochester DF. Does respiratory muscle rest relieve fatigue or incipient fatigue? *Am Rev Respir Dis* 1988;138:516-517.
- Swartz MA, Marino PL. Diaphragmatic strength during weaning from mechanical ventilation. *Chest* 1985;88:736-739.
- Decramer J, Lacquet LM, Fagard R, et al. Corticosteroids contribute to muscle weakness in chronic airflow obstruction. *Am J Respir Crit Care Med* 1994;150:11-16.
- Gallagher CG. Respiratory steroid myopathy. *Am J Respir Crit Care Med* 1994;150:4-6.
- Lee C. Intensive care unit neuromuscular syndrome? *Anesthesiology* 1995;83:237-240.
- Rossi A, Gottfried SB, Zocchi L, Higgs BD, Lennox S, Calverley PM, et al. Measurement of static compliance of the total respiratory system in patients with acute respiratory failure during mechanical ventilation: the effect of intrinsic positive end-expiratory pressure. *Am Rev Respir Dis* 1985;131(5):672-677.
- Hilberman M, Kamm B, Lamy M, et al. An analysis of potential physiological predictors of respiratory adequacy following cardiac surgery. *J Thorac Cardiovasc Surg* 1976;71:711-720.
- Jabour ER, Rabil DM, Truweit JD, Rochester DF. Evaluation of a new weaning index based on ventilatory endurance and the efficiency of gas exchange. *Am Rev Respir Dis* 1991;144:531-537.
- Sassoon CS, Mahutte CK. Airway occlusion pressure and breathing pattern as predictors of weaning outcome. *Am Rev Respir Dis* 1993;148:860-866.
- Kline JL, Zimnicki GL, Antonenko DR, Bander JJ. The use of calculated relative inspiratory effort as a predictor of outcome in mechanical ventilation weaning trials. *Respir Care* 1987;32:870-876.
- el-Khatib MF, Baumeister B, Smith PG, Chatburn RL, Blumer JL. Inspiratory pressure/maximal inspiratory pressure ($P_{I_{max}}$): does it predict successful extubation in critically ill infants and children?. *Int Care Med* 1996;22:264-268.
- Khan N, Brown A, Venkataraman ST. Predictors of extubation success and failure in mechanically ventilated infants and children. *Crit Care Med* 1996;24:1568-1579.
- Leitch EA, Moran JL, Grealy B. Weaning and extubation in the intensive care unit: clinical or index-driven approach? *Int Care Med* 1996;22:752-759.
- Mergoni M, Costa A, Primavera S, Salvadori A, Saccani A, Zuccoli P. [Assessment of various new predictive parameters of the outcome of mechanical ventilation weaning]. Article in Italian. *Minerva Anestesiol* 1996;62:153-164.
- Gologorskii VA, Gelfand BR, Stamov VI, Lapshina II, Nistratov SL. [Cessation of prolonged artificial ventilation of the lungs and transition to spontaneous respiration of surgical patients]. Article in Russian. *Anesteziol Reanimatol* 1997;1:4-10.

43. Krieger BP, Isber J, Breitenbucher A, Throop G, Ershowsky P. Serial measurements of the rapid-shallow-breathing index as a predictor of weaning outcome in elderly medical patients. *Chest* 1997; 112:1029–1034.
44. Krieger BP, Ershowsky PF, Becker DA, Gazeroglu HB. Evaluation of conventional criteria for predicting successful weaning from mechanical ventilatory support in elderly patients. *Crit Care Med* 1989; 17:858–861.
45. Ashutosh K, Lee H, Mohan CK, Ranka S, Mehrotra K, Alexander C. Prediction criteria for successful weaning from respiratory support: statistical and connectionist analyses. *Crit Care Med* 1992;20: 1295–1301. [published erratum appears in *Crit Care Med* 1994 Jan;22(1):183.]
46. Sahn SA, Lakshminarayan S. Bedside criteria for discontinuation of mechanical ventilation. *Chest* 1973;63:1002–1005.
47. Gandia F, Blanco J. Evaluation of indexes predicting the outcome of ventilator weaning and value of adding supplemental inspiratory load. *Int Care Med* 1992;18:327–333.
48. Farias JA, Alia I, Esteban A, Golubicki AN, Olazarri FA. Weaning from mechanical ventilation in pediatric intensive care patients. *Int Care Med* 1998;24:1070–1075.
49. Peters RM, Brimm JE, Utley JR. Predicting the need for prolonged ventilatory support in adult cardiac patients. *J Thoracic Cardiovasc Surg* 1979;77:175–182.
50. Cohen CA, Zigelbaum G, Gross D, Roussos C, Macklem PT. Clinical manifestations of inspiratory muscle fatigue. *Am J Med* 1982; 73:308–316.
51. Esau SA, Bellemare F, Grassino A, Permutt S, Roussos C, Pardy RL. Changes in relaxation rate with diaphragmatic fatigue in humans. *J Appl Physiol* 1983;54:1353–1360.
52. Jubran A, Tobin MJ. Pathophysiological basis of acute respiratory distress in patients who fail a trial of weaning from mechanical ventilation. *Am J Respir Crit Care Med* 1997;155:906–915.
53. Mier-Jedrzejowicz A, Brophy C, Moxham J, Green M. Assessment of diaphragm weakness. *Am Rev Respir Dis* 1988;137:877–883.
54. Tobin MJ. 1999 Donald F Egan Scientific Lecture: Weaning from mechanical ventilation: what have we learned? *Respir Care* 2000; 45:417–431.
55. De Troyer A, Estenne M. Coordination between rib cage muscles and diaphragm during quiet breathing in humans. *J Appl Physiol* 1984;57:899–906.
56. Gallagher CG, Im Hof V, Younes M. Effect of inspiratory muscle fatigue on breathing pattern. *J Appl Physiol* 1985;59:1152–1158.
57. Gilbert R, Auchincloss JH, Peppi D, Ashutosh K. The first few hours off a respirator. *Chest* 1974;65:152–157.
58. Sharp JT, Goldberg NB, Druz WS, Fishman HC, Danon J. Thoracoabdominal motion in chronic obstructive pulmonary disease. *Am Rev Respir Dis* 1977;115:47–56.
59. Tobin MJ, Perez W, Guenther SM, Lodato RF, Dantzker DR. Does rib cage-abdominal paradox signify respiratory muscle fatigue? *J Appl Physiol* 1987;63:851–860.
60. Tobin MJ, Guenther SM, Perez W, et al. Konno-Mead analysis of rib cage-abdominal motion during successful and unsuccessful trials of weaning from mechanical ventilation. *Am Rev Respir Dis* 1987;135:1320–1328.
61. Aubier M, Murciano D, Lecocguic Y, Viires N, Pariente R. Bilateral phrenic stimulation: a simple technique to assess diaphragmatic fatigue in humans. *J Appl Physiol* 1985;58:58–64.
62. Dunn WF, Nelson SB, Hubmayr RD. The control of breathing during weaning from mechanical ventilation. *Chest* 1991;100:754–761.
63. Field S, Kelly SM, Macklem PT. The oxygen cost of breathing in patients with cardiorespiratory disease. *Am Rev Respir Dis* 1982; 126:9–13.
64. Hubmayr RD, Loosbrock LM, Gillespie DJ, Rodarte JR. Oxygen uptake during weaning from mechanical ventilation. *Chest* 1988; 94:1148–1155.
65. Murciano D, Boczkowski J, Lecocguic Y, Milic-Emili J, Pariente R, Aubier M. Tracheal occlusion pressure: a simple index to monitor respiratory muscle fatigue during acute respiratory failure in patients with chronic obstructive pulmonary disease. *Ann Intern Med* 1988;108:800–805.
66. Stroetz RW, Hubmayr RD. Tidal volume maintenance during weaning with pressure support. *Am J Respir Crit Care Med* 1995;152: 1034–1040.
67. Gay PC, Rodarte JR, Hubmayr RD. The effects of positive expiratory pressure on isovolume flow and dynamic hyperinflation in patients receiving mechanical ventilation. *Am Rev Respir Dis* 1989; 139:621–626.
68. Parthasarathy S, Jubran A, Tobin MJ. Cycling of inspiratory and expiratory muscle groups with the ventilator in airflow limitation. *Am J Respir Crit Care Med* 1998;158:1471–1478.
69. Pepe PE, Marini JJ. Occult positive end-expiratory pressure in mechanically ventilated patients with airflow obstruction: the auto-PEEP effect. *Am Rev Respir Dis* 1982;126:166–170.
70. Reinoso MA, Gracey DR, Hubmayr RD. Interrupter mechanics of patients admitted to a chronic ventilator dependency unit. *Am Rev Respir Dis* 1993;148:127–131.
71. Sassoon CSH, Te TT, Mahutte K, Light RW. Airway occlusion pressure: an important indicator for successful weaning in patients with chronic obstructive pulmonary disease. *Am Rev Respir Dis* 1987;135:107–113.
72. Gandevia SC. Neural control in human muscle fatigue: changes in muscle afferents, motoneurons and motor cortical drive. *Acta Physiologica Scandinavica* 1998;162(2):275–283.
73. Yang KL, Tobin MJ. A prospective study of indexes predicting the outcome of trials of weaning from mechanical ventilation. *N Engl J Med* 1991;324:1445–1450.
74. Pingleton SK, Harmon GS. Nutritional management in acute respiratory failure. *JAMA* 1987;257:3094–3099.
75. Lewis MI, Sieck GC, Fournier M, et al. The effect of nutritional deprivation of diaphragm contractility and muscle fiber size. *J Appl Physiol* 1986;60:596–603.
76. Doekel RC, Zwillich C, Scoggin CH, Kryger M, Weil JV. Clinical semi-starvation: depression of hypoxic ventilatory response. *N Engl J Med* 1976;295:358–361.
77. Bassili HR, Deitel M. Effect of nutritional support on weaning patients off mechanical ventilators. *J Parenter Enteral Nutr* 1981; 5(2):161–163.
78. Larca L, Greenbaum DM. Effectiveness of intensive nutritional regimes in patients who fail to wean from mechanical ventilation. *Crit Care Med* 1982;10:297–300.
79. Aubier M, Murciano D, Lecocguic Y, et al. Effects of hypophosphatemia on diaphragmatic contractility in patients with acute respiratory failure. *N Engl J Med* 1985;313:420–424.
80. Aubier M, Viires N, Piquet J, et al. Effects of hypocalcemia on diaphragmatic strength generation. *J Appl Physiol* 1985;58:2054–2061.
81. Cerra F. Hypermetabolism, organ failure and metabolic support. *Surgery* 1987;101:1–14.
82. Dhingra S, Solven F, Wilson A, McCarthy DS. Hypomagnesemia and respiratory muscle power. *Am Rev Respir Dis* 1984;129:497–498.
83. Zwillich CW, Pierson DJ, Hofeldt FD, et al. Ventilatory control in myxedema and hypothyroidism. *N Engl J Med* 1975;292:662–665.
84. Siafakas NM, Salesiotou V, Filaditaki V, et al. Respiratory muscle strength in hypothyroidism. *Chest* 1992;102:189–194.

85. Mohsenifar Z, Hay A, Hay J, Lewis MI, Koerner SK. Gastric intramural pH as a predictor of success or failure in weaning patients from mechanical ventilation. *Ann Int Med* 1993;119:794–798.
86. Bouachour G, Guiraud MP, Gouello JP, Roy PM, Alquier P. Gastric intramucosal pH: an indicator of weaning outcome from mechanical ventilation in COPD patients. *Eur Respir J* 1996;9:1868–1873.
87. Bando K, Sun K, Binford RS, et al. Determinants of longer duration of endotracheal intubation after adult cardiac operations. *Ann Thor Surg* 1997;63:1026–1033.
88. Artigas A, Bernard GR, Carlet J, et al. The American European Consensus Conference on ARDS, Part 2. Ventilatory, pharmacologic, supportive therapy, study design strategies, and issues related to recovery and remodeling. *Am J Respir Crit Care Med* 1998;157:1332–1347.
89. Jubran A, Mathru M, Dries D, Tobin MJ. Continuous recordings of mixed venous oxygen saturation during weaning from mechanical ventilation and the ramifications thereof. *Am J Respir Crit Care Med* 1998;158:1763–1769.
90. Lemaire F, Teboul J-L, Cinotti L, Giotto G, Abrouk F, Steg G, et al. Acute left ventricular dysfunction during unsuccessful weaning from mechanical ventilation. *Anesthesiology* 1988;69:171–179.
91. Epstein SK. Etiology of extubation failure and the predictive value of the rapid shallow breathing index. *Am J Respir Crit Care Med* 1995;152:545–549.
92. Richard CH, Teboul JL, Archambaud F, et al. Left ventricular function during weaning of patients with chronic obstructive pulmonary disease. *Intensive Care Med* 1994;20:181–186.
93. Chatila W, Ani S, Guaglianone D, et al. Cardiac ischemia during weaning from mechanical ventilation. *Chest* 1996;109:1577–1583.
94. Nett LM, Morganroth M, Petty TL. Weaning from mechanical ventilation: a perspective and review of techniques. In: Bone RC, editor. *Critical care: a comprehensive approach*. Park Ridge IL: Am Coll Chest Phys; 1984:171–188.
95. Cooper KR, Phillips BA. Effect of short-term sleep loss on breathing. *J Appl Physiol* 1982;53:855–858.
96. Spengler CM, Shea SA. Sleep deprivation per se does not decrease the hypercapnic ventilatory response in humans. *Am J Respir Crit Care Med* 2000;161:1124–1128.
97. LaRiccica PJ, Katz RH, Peters JW, et al. Biofeedback and hypnosis in weaning from mechanical ventilators. *Chest* 1985;87:267–269.
98. Holiday JE, Hyers TM. The reduction of weaning time from mechanical ventilation using tidal volume and relaxation feedback. *Am Rev Respir Dis* 1990;141:1214–1220.
99. Torres A, Serra-Batlles J, Ros E, et al. Pulmonary aspiration of gastric contents in patients receiving mechanical ventilation: the effect of body position. *Ann Int Med* 1992;116:540–543.
100. Epstein SK, Ciubotaru RL, Wong JB. Effect of failed extubation on the outcome of mechanical ventilation. *Chest* 1997;112:186–192.
101. Esteban A, Alia I, Gordo F, Fernandez R, Solsona JF, Vallverdu I, et al. Extubation outcome after spontaneous breathing trials with T-tube or pressure support ventilation. The Spanish Lung Failure Collaborative Group. *Am J Respir Crit Care Med* 1997;156(2 pt 1):459–465. [published erratum appears in *Am J Respir Crit Care Med* 1997;156(6):2028.]
102. Esteban A, Alia I, Tobin MJ, Gil A, Gordo F, Vallverdu I, et al. Effect of spontaneous breathing trial duration on outcome of attempts to discontinue mechanical ventilation. Spanish Lung Failure Collaborative Group. *Am J Respir Crit Care Med* 1999;159(2):512–518.
103. Vallverdu I, Calaf N, Subirana M, Net A, Benito S, Mancebo J. Clinical characteristics, respiratory functional parameters, and outcome of a two-hour T-piece trial in patients weaning from mechanical ventilation. *Am J Respir Crit Care Med* 1998;158:1855–1862.
104. Ely EW, Baker AM, Evans GW, Haponik EF. The prognostic significance of passing a daily screen of breathing spontaneously. *N Engl J Med* 1999;25:581–587.
105. Fagon JY, Chastre J, Hance AJ, Montravers P, Novara A, Gibert C. Nosocomial pneumonia in ventilated patients: a cohort study evaluating attributable mortality and hospital stay. *Am J Med* 1993;94:281–288.
106. Epstein SK, Ciubotaru RL. Independent effects of etiology of failure and time to reintubation on outcome for patients failing extubation. *Am J Respir Crit Care Med* 1998;158:489–493.
107. Esteban A, Frutos F, Tobin MJ, Alia I, Solsona JF, Valverdu I, et al. A comparison of four methods of weaning patients from mechanical ventilation. Spanish Lung Failure Collaborative Group. *N Engl J Med* 1995;332:345–350.
108. Ely EW, Baker AM, Dunagan DP, Burke HL, Smith AC, Kelly PT, et al. Effect on the duration of mechanical ventilation of identifying patients capable of breathing spontaneously. *N Engl J Med* 1996;335:1864–1869.
109. Brochard L, Rauss A, Benito S, Conti G, Mancebo J, Rekik N, et al. Comparison of three methods of gradual withdrawal from ventilatory support during weaning from mechanical ventilation. *Am J Respir Crit Care Med* 1994;150:896–903.
110. Dojat M, Harf A, Touchard D, Laforest M, Lemaire F, Brochard L. Evaluation of a knowledge-based system providing ventilatory management and decision for extubation. *Am J Respir Crit Care Med* 1996;153:997–1004.
111. Yang KL. Reproducibility of weaning parameters: a need for standardization. *Chest* 1992;102:1829–1832.
112. Multz AS, Aldrich TK, Prezant DJ, Karpel JP, Hendler JM. P_{1max} is not a reliable test of inspiratory muscle strength in mechanically ventilated patients. *Am Rev Respir Dis* 1990;142:529–532.
113. Burns SM, Clochesy JM, Goodnough-Hanneman SK, et al. Weaning from long-term mechanical ventilation. *Am J Crit Care Med* 1995;4:4–22.
114. Goodnough-Hanneman SK, Ingersoll GL, Knebel AR, et al. Weaning from short-term mechanical ventilation: a review. *Am J Crit Care Med* 1994;3:421–443.
115. Jaeschke R, Guyatt G, Sackett D, et al, for the Evidence-Based Medicine Working Group. Users' guide to the medical literature: how to use an article about a diagnostic test. B. What are the results and will they help me in caring for my patients? *JAMA* 1994;271:703–707.
116. Jubran A, Parthasarathy S. Hypercapnic respiratory failure during weaning: neuromuscular capacity vs muscle loads. *Respir Care Clin N Am* 2000;6:385–406.
117. Capdevila X, Perrigault PF, Ramonatxo M, Roustan JP, Peray P, d'Athis F, Prefaut C. Changes in breathing pattern and respiratory muscle performance parameters during difficult weaning. *Crit Care Med* 1998;26:79–87.
118. Vassilakopoulos T, Zakynthinos S, Roussos C. The tension-time index and the frequency/tidal volume ratio are the major pathophysiological determinants of weaning failure and success. *Am J Respir Crit Care Med* 1998;158:378–385.
119. Kollef MH, Shapiro SD, Silver P, Prentice D, Sauer S, Ahrens TS, et al. A randomized, controlled trial of protocol-directed versus physician-directed weaning from mechanical ventilation. *Crit Care Med* 1997;25:567–574.
120. Chopin C, Chambrin MC, Mangalaboyi J, Lestavel P, Fourrier F. Carbon dioxide mandatory ventilation (CO_2MV): a new method for weaning from mechanical ventilation—description and compar-

- tive clinical study with I.M.V. and T. tube method in COPD patient. *Int J Clin Monit Comput* 1989;6:11–19.
121. Stroetz RW, Hubmayr RD. Tidal volume maintenance during weaning with pressure support. *Am J Respir Crit Care Med* 1995;152:1034–1040.
 122. Engoren M, Buderer NF, Zacharias A, Habib RH. Variables predicting reintubation after cardiac surgical procedures. *Ann Thoracic Surg* 1999;67:661–665.
 123. Rady MY, Ryan T. Perioperative predictors of extubation failure and the effect on clinical outcome after cardiac surgery. *Crit Care Med* 1999;27:340–347.
 124. London MJ, Shroyer AL, Coll JR, MaWhinney S, Fullerton DA, Hammermeister KE, Grover FL. Early extubation following cardiac surgery in a veterans population. *Anesthesiology* 1998;88:1447–1458.
 125. deHaven CB, Kirton OC, Morgan JP, Hart AM, Shatz DV, Civetta JM. Breathing measurements reduces false negative classification of tachypneic pre-extubation trial failures. *Crit Care Med* 1996;24:976–980.
 126. Laghi F, D'Alfonso N, Tobin MJ. Pattern of recovery from diaphragmatic fatigue over 24 hours. *J Appl Physiol* 1995;79:539–546.
 127. Reid WD, Huang J, Bryson S, Walker DC, Belcastro AN. Diaphragm injury and myofibrillar structure induced by resistive loading. *J Appl Physiol* 1994;76:176–184.
 128. Cohen CA, Zagalbaum G, Gross D, Roussos C, Macklem PT. Clinical manifestations of inspiratory muscle fatigue. *Am J Med* 1982;73:308–316.
 129. Feeley TW, Saumarez R, Klick JM, McNabb TG, Skillman JJ. Positive end-expiratory pressure in weaning patients from controlled ventilation: a prospective randomized trial. *Lancet* 1975;2(7938):725–729.
 130. Jones DP, Byrne P, Morgan C, Fraser I, Hyland R. Positive end-expiratory pressure vs T-piece: extubation after mechanical ventilation. *Chest* 1991;100:1655–1659.
 131. Bailey CR, Jones RM, Kelleher AA. The role of continuous positive airway pressure during weaning from mechanical ventilation in cardiac surgical patients. *Anaesthesia* 1995;50:677–681.
 132. Petrof BJ, Legare M, Goldberg P, et al. Continuous positive airway pressure reduces work of breathing and dyspnea during weaning from mechanical ventilation in severe chronic obstructive pulmonary disease. *Am Rev Respir Dis* 1990;141:281–289.
 133. Smith TC, Marini JJ. Impact of PEEP on lung mechanics and work of breathing in severe airflow obstruction. *J Appl Physiol* 1988;65:1488–1497.
 134. Fisher MM, Raper RF. The “cuff leak” test for extubation. *Anaesthesia* 1992;47:10–12.
 135. Miller R, Cole R. Association between reduced cuff leak volume and postextubation stridor. *Chest* 1996;110:1035–1040.
 136. Engoren M. Evaluation of the cuff leak test in cardiac surgery patients. *Chest* 1999;116:1029–1031.
 137. Coplin WM, Pierson DJ, Cooley KD, Newell DW, Reubenfeld GD. Implications of extubation delay in brain injured patients meeting standard weaning criteria. *Am J Respir Crit Care Med* 2000;161:1530–1536.
 138. Capdevila XJ, Perrigault PF, Perey PJ, Roustan JP, d'Athis F. Occlusion pressure and its ratio to maximum inspiratory pressure are useful predictors for successful extubation following T-piece weaning trial. *Chest* 1995;108:482–489.
 139. Bach J, Saporto L. Criteria for extubation and tracheostomy tube removal for patients with ventilatory failure. *Chest* 1996;110:1566–1571.
 140. Travaline JM, Sudershan S, Criner GJ. Recovery of Pdi twitch following the induction of diaphragm fatigue in normal subjects. *Am J Respir Crit Care Med* 1997;156:1562–1566.
 141. Jounieaux V, Duran A, Levi-Valensi P. Synchronized intermittent mandatory ventilation with and without pressure support ventilation in weaning patients with COPD from mechanical ventilation. *Chest* 1994;105:1204–1210.
 142. Esen F, Denkel T, Telci L, Kesecioglu J, Tutuncu AS, Akpir K, Lachmann B. Comparison of pressure support ventilation (PSV) and intermittent mandatory ventilation (IMV) during weaning in patients with acute respiratory failure. *Adv Exper Med Biol* 1992;317:371–376.
 143. Hess D, Branson RD. Ventilators and weaning modes. *Respir Care Clin N Am* 2000;6:407.
 144. Straus C, Louis B, Isabey D, et al. Contribution of the endotracheal tube and the upper airway to breathing workload. *Am J Respir Crit Care Med* 1998;157:23–30.
 145. Iotti GA, Olivei MC, Palo A, et al. Unfavorable mechanical effects of heat and moisture exchangers in ventilated patients. *Int Care Med* 1997;23:399–405.
 146. LeBourdelle G, Mier L, Fiquet B, et al. Comparison of the effects of heat and moisture exchangers and heated humidifiers on ventilation and gas exchange during weaning trials from mechanical ventilation. *Chest* 1996;110:1294–1298.
 147. Sassoon CSH. Mechanical ventilator design and function: the trigger variable. *Respir Care* 1992;37:1056–1069.
 148. MacIntyre NR, Ho L. Effects of initial flow rate and breath termination criteria on pressure support ventilation. *Chest* 1991;99:134–138.
 149. MacIntyre NR, et al. Patient-ventilator flow dyssynchrony. flow limited versus pressure limited breaths. *Crit Care Med* 1997;25:1671.
 150. Branson RD, Campbell RS, Davis K, et al. Altering flowrate during maximum pressure support ventilation (PSV_{max}): Effects on cardiorespiratory function. *Respir Care* 1990;35:1056–1064.
 151. Imsand C, Feihl F, Perret C, Fitting JW. Regulation of inspiratory neuromuscular output during synchronized intermittent mechanical ventilation. *Anesthesiology* 1994;80:13–22.
 152. Marini JJ, Smith TC, Lamb VJ. External work output and force generation during synchronized intermittent mandatory ventilation: effect of machine assistance on breathing effort. *Am Rev Respir Dis* 1988;138:1169–1179.
 153. Parthasarathy S, Jubran A, Tobin MJ. Cycling of inspiratory and expiratory muscle groups with the ventilator in airflow limitation. *Am J Respir Crit Care Med* 1998;158:1471–1478.
 154. Jubran A, van de Graaff WB, Tobin MJ. Variability of patient-ventilator interaction with pressure support ventilation in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1995;152:129–136.
 155. Branson RD, MacIntyre NR. Dual-control modes of mechanical ventilation. *Respir Care* 1996;41:294–305.
 156. Linton DM, Potgieter PD, Davis S, et al. Automatic weaning from mechanical ventilation using an adaptive lung controller. *Chest* 1994;105:1843–1850.
 157. Dojat M, Harf A, Touchard D, LeMaire F, Brochard L. Clinical evaluation of a computer controlled pressure support mode. *Am J Respir Crit Care Med* 2000;161:1161–1166.
 158. Davis S, Potgieter PD, Linton DM. Mandatory minute volume weaning in patients with pulmonary pathology. *Anaesth Int Care* 1989;17:170–174.
 159. Girault C, Daudenthun I, Chevron V, Tamion F, Leroy J, Bonmarchand G. Noninvasive ventilation as a systematic extubation and weaning technique in acute-on-chronic respiratory failure: a prospective, randomized controlled study. *Am J Respir Crit Care Med* 1999;160:86–92.
 160. Nava S, Ambrosino N, Clini E, Prato M, Orlando G, Vitacca M, et al. Noninvasive mechanical ventilation in the weaning of patients

- with respiratory failure due to chronic obstructive pulmonary disease: a randomized, controlled trial. *Ann Intern Med* 1998;128:721–728.
161. Cheng DC, Karski J, Peniston C, Asokumar B, Raveendran G, Carroll J, et al. Morbidity outcome in early versus conventional tracheal extubation after coronary artery bypass grafting: a prospective randomized controlled trial. *J Thor Cardiovasc Surg* 1996;112:755–764.
 162. Berry PD, Thomas SD, Mahon SP, Jackson M, Fox MA, Fabri B, et al. Myocardial ischaemia after coronary artery bypass grafting: early vs. late extubation. *Br J Anaesth* 1998;80:20–25.
 163. Michalopoulos A, Nikolaidis A, Antzaka C, Deliyanni M, Smirli A, Geroulanos S, Papadimitriou L. Change in anaesthesia practice and postoperative sedation shortens ICU and hospital length of stay following coronary artery bypass surgery. *Respir Med* 1998;92:1066–1070.
 164. Silbert BS, Santamaria JD, O'Brien JL, Blyth CM, Kelly WJ, Molnar RR. Early extubation following coronary artery bypass surgery: a prospective randomized controlled trial. The Fast Track Cardiac Care Team. *Chest* 1998;113:1481–1488.
 165. Engoren MC, Kraras C, Garzia F. Propofol-based versus fentanyl-isoflurane-based anaesthesia for cardiac surgery. *J Cardiothorac Vasc Anesth* 1998;12:177–181.
 166. Shackford SR, Virgilio RW, Peters RM. Early extubation versus prophylactic ventilation in the high risk patient: a comparison of postoperative management in the prevention of respiratory complications. *Anesth Analg* 1981;60:76–80.
 167. Tempe D, Cooper A, Mohan JC, Nigam M, Tomar AS, Ramesh K, et al. Closed mitral valvotomy and elective ventilation in the postoperative period: effect of mild hypercarbia on right ventricular function. *J Cardiothorac Vasc Anesth* 1995;9:552–557.
 168. Quasha AL, Loeber N, Freeley TW, Ulyot DH, Roizen MF. Postoperative respiratory care: a controlled trial of early and late extubation following coronary-artery bypass grafting. *Anesthesiology* 1980;52:135–141.
 169. Reyes A, Vega G, Blancas R, Morato B, Moreno JL, Torrecilla C, Cereijo E. Early vs conventional extubation after cardiac surgery with cardiopulmonary bypass. *Chest* 1997;112:193–201.
 170. Dumas A, Dupuis GH, Searle N, et al. Early versus late extubation after coronary artery bypass grafting: effects on cognitive function. *J Cardiothorac Vasc Anesth* 1999;13:130–135.
 171. Stoller JK, Mascha EJ, Kester L, Haney DJ. Randomized controlled trial of physician-directed versus respiratory therapy consult service-directed respiratory care to adult non-ICU inpatients. *Am J Respir Crit Care Med* 1998;158:1068–1075.
 172. Marelich GP, Murin S, Battistella F, Inciardi J, Vierra T, Roby M. Protocol weaning of mechanical ventilation in medical and surgical patients by respiratory therapists and nurses: effect on weaning time and incidence of ventilator-associated pneumonia. *Chest* 2000;118:459–467.
 173. Brook AD, Ahrens TS, Schaiff R, et al. Effect of a nursing-implemented sedation protocol on the duration of mechanical ventilation. *Crit Care Med* 1999;27(12):2609–2615.
 174. Kress JP, Pohlman AS, O'Connor ME, Hall JB. Daily interruption of sedative infusions in critically ill patients undergoing mechanical ventilation. *N Engl J Med*. 2000;342:1471–1477.
 175. Ely EW, Namen AM, Tatter S, Lucia MA, Smith AC, Landry S, et al. Impact of a ventilator weaning protocol in neurosurgical patients: a randomized, controlled trial (abstract). *Am J Respir Crit Care Med* 1999;159:A371.
 176. Collin GR, Atkinson N, Furrow K, Trenis T, Keeley RC, Sedovy JD, et al. Decreasing the duration of mechanical ventilation through the use of a ventilation management protocol (abstract). *Crit Care Med* 1999;27:A110.
 177. Ely EW, Bennett PA, Bowton DL, Murphy SM, Haponik EF. Large scale implementation of a respiratory therapist-driven protocol for ventilator weaning. *Am J Respir Crit Care Med* 1999;159:439–446.
 178. Fridkin SK, Pear SM, Williamson TH, Galgiani JN, Jarvis WR. The role of understaffing in central venous catheter-associated bloodstream infections. *Infect Control Hosp Epidemiol* 1996;17:150–158.
 179. Archibald LK, Manning ML, Bell LM, Banedee S, Jarvis VTR. Patient density, nurse-to-patient ratio and nosocomial infection risk in a pediatric cardiac intensive care unit. *Pediatr Infect Dis* 1997;16:1046–1048.
 180. Thorens JB, Kaelin RM, Jolliet P, Chevrolet J-C. Influence of the quality of nursing on the duration of weaning from mechanical ventilation in patients with chronic obstructive pulmonary disease. *Crit Care Med* 1995;23:1807–1815.
 181. Plummer AL, Gracey DR. Consensus conference on artificial airways in patients receiving mechanical ventilation. *Chest* 1989;96:178–184.
 182. Stauffer JL, Olson DE, Petty TL. Complications and consequences of endotracheal intubation and tracheostomy: a prospective study of 150 critically ill adult patients. *Am J Med* 1981;70:65–76.
 183. Astrachan DI, Kirchner JC, Goodwin WJJ. Prolonged intubation vs tracheostomy: complications, practical and psychological considerations. *Laryngoscope* 1988;98:1165–1169.
 184. Stock MC, Woodward CG, Shapiro BA, et al. Perioperative complications of elective tracheostomy in critically ill patients. *Crit Care Med* 1986;14:861–863.
 185. Boyd SW, Benzel EC. The role of early tracheostomy in the management of the neurosurgical patient. *Laryngoscope* 1992;102:559–562.
 186. Heffner JE. Tracheostomy: indications and timing. *Respir Care* 1999;44:807–815.
 187. Heffner JE. Timing of tracheostomy in mechanically ventilated patients. *Am Rev Respir Dis* 1993;147:768–771.
 188. Colice GL, Stukel TA, Dain B. Laryngeal complications of prolonged intubation. *Chest* 1989;96:877–884.
 189. Whited RE. A prospective study of laryngotracheal sequelae in long-term intubation. *Laryngoscope* 1984;94:367–377.
 190. Kollef MH, Levy NT, Ahrens TS, et al. The use of continuous intravenous sedation is associated with prolongation of mechanical ventilation. *Chest* 1998;114:541–548.
 191. Bersten AD, Rutten AJ, Vedig AE, et al. Additional work of breathing imposed by endotracheal tubes, breathing circuits, and intensive care ventilators. *Crit Care Med* 1989;17:671–677.
 192. Habib M. Physiological implications of artificial airways. *Chest* 1989;96:180–184.
 193. Yung MW, Snowdon SL. Respiratory resistance of tracheostomy tubes. *Arch Otolaryngol* 1984;110:591–595.
 194. Wright PE, Marini JJ, Bernard GR. In vitro versus in vivo comparison of endotracheal tube airflow resistance. *Am Rev Respir Dis* 1989;140:10–16.
 195. Diehl JL, El Atrous S, Touchard D, et al. Changes in the work of breathing induced by tracheostomy in ventilator-dependent patients. *Am J Respir Crit Care Med* 1999;159:383–388.
 196. Davis K, Jr, Campbell RS, Johannigman JA, et al. Changes in respiratory mechanics after tracheostomy. *Arch Surg* 1999;134:59–62.
 197. Lin MC, Huang CC, Yang CT, et al. Pulmonary mechanics in patients with prolonged mechanical ventilation requiring tracheostomy. *Anaesth Intensive Care* 1999;27:581–585.
 198. Rodriguez JL, Steinberg SM, Luchetti FA, et al. Early tracheostomy for primary airway management in the surgical critical care setting. *Surgery* 1990;108:655–659.

199. Dunham MC, LaMonica C. Prolonged tracheal intubation in the trauma patient. *Trauma* 1984;24:120–124.
200. Sugeran HJ, Wolfe L, Pasquale MD, et al. Multicenter, randomized, prospective trial of early tracheostomy. *J Trauma* 1997;43:741–747.
201. Lesnik I, Rappaport W, Fulginiti J, et al. The role of early tracheostomy in blunt, multiple organ trauma. *Am Surg* 1992;58:346–349.
202. Blot F, Guiguet M, Antoun S, et al. Early tracheotomy in neutropenic, mechanically ventilated patients: rationale and results of a pilot study. *Support Care Cancer* 1995;3:291–296.
203. Koh WY, Lew TWK, Chin NM, et al. Tracheostomy in a neuro-intensive care setting: indications and timing. *Anaesth Intensive Care* 1997;25:365–368.
204. El-Naggar M, Sadagopan S, Levine H, et al. Factors influencing choice between tracheostomy and prolonged translaryngeal intubation in acute respiratory failure: a prospective study. *Anesth Analg* 1976;55:195–201.
205. Maziak DE, Meade MO, Todd TRJ. The timing of tracheotomy: a systematic review. *Chest* 1998;114:605–609.
206. Seneff MG, Zimmerman JE, Knaus WA, et al. Predicting the duration of mechanical ventilation: the importance of disease and patient characteristics. *Chest* 1996;110:469–479.
207. Kurek CK, Cohen IL, Lambrinos J, et al. Clinical and economic outcome of patients undergoing tracheostomy for prolonged mechanical ventilation in New York State during 1993: analysis of 6,353 cases under diagnostic related group 483. *Crit Care Med* 1997;25:983–988.
208. Scheinhorn DJ, Chao DC, Steam-Hassenpflug MA, et al. Post-ICU mechanical ventilation. Treatment of 1,123 patients at a regional weaning center. *Chest* 1997;111:1654.
209. Scheinhorn DJ, Artinian BA, Catlin JL. Weaning from prolonged mechanical ventilation: the experience at a regional weaning center. *Chest* 1994;105:534.
210. Scheinhorn DJ, Hassenpflug M, Artinian B. Predictors of weaning after six weeks of mechanical ventilation. *Chest* 1995;107:500.
211. Gracey DR, Hardy DC, Naessens JM, et al. The mayo ventilator-dependent rehabilitation unit: a 5-year experience. *Mayo Clin Proc* 1997;72:13.
212. Elpern EH, Larson R, Douglass P, et al. Long-term outcomes for elderly survivors of prolonged ventilator assistance. *Chest* 1989;96:1120–1124.
213. Indihar FJ. A 10-year report of patients in a prolonged respiratory care unit. *Minn Med* 1991;74:23–27.
214. Latriano B, McCauley P, Astiz ME, et al. Non-ICU care of hemodynamically stable mechanically ventilated patients. *Chest* 1996;109:1591.
215. Petrak RA, Nicholson KI, Brofman JD. Clinical outcomes prediction based on demographic data at a regional chronic ventilator-dependent unit. Program and abstracts of “Weaning ’96,” Weaning from prolonged mechanical ventilation. Palm Springs, 1996.
216. Clark RL, Theiss D. Prolonged mechanical ventilation weaning, the experience at an extended critical care regional weaning center (abstract). *Am J Respir Crit Care Med* 1997;155:A410.
217. Bagley PH, Cooney E. A community-based regional ventilator weaning unit. *Chest* 1997;111:1024–1029.
218. Carson SS, Bach PB, Brzozowski L, et al. Outcomes after long term acute care: an analysis of 133 mechanically ventilated patients. *Am J Respir Crit Care Med* 1999;159:1568–1573.
219. Dasgupta A, Rice R, Mascha E, et al. Four-year experience with a unit for long-term ventilation (respiratory special care unit) at the Cleveland Clinic Foundation. *Chest* 1999;116:447–455.
220. Gracey DR, Hardy DC, Koenig GE. Cost effective alternatives to intensive unit care: experience in the Mayo Rochester Chronic Ventilator Dependent Unit. *Mayo Clin Proc* 2000;75:445–449.
221. Burns SM, Marshall M, Burns JE, et al. Design, testing, and results of an outcomes-managed approach to patients requiring prolonged mechanical ventilation. *Am J Crit Care* 1998;7:45–57.
222. Ruby EB, Dakly BJ, Douglas S, Montenegro HD, Song R, Dyer, MA. Patient outcomes for the chronically critically ill: special care vs intensive care unit. *Nurs Res* 1995;44:324–331.
223. Scheinhorn DJ, Chao DC, Hassenpflug MS, et al. Post-ICU mechanical ventilation: results of a therapist-implemented patient-specific (TIPS) weaning protocol (abstract). *Am J Respir Crit Care Med* 1999;159:A373.
224. Scheinhorn DJ, Chao DC, Hassenpflug MS. Approach to patients with long-term weaning failure. *Respir Care Clin N Am* 2000;6:437–461.