

Nos. 20-1530, 20-1531, 20-1778, and 20-1780

IN THE
Supreme Court of the United States

STATE OF WEST VIRGINIA, ET AL., *Petitioners,*

v.

U.S. ENVIRONMENTAL PROTECTION AGENCY, ET AL.,
Respondents.

On Writ of Certiorari to the United States Court of
Appeals for the District of Columbia Circuit

**BRIEF OF AMICI CURIAE AMERICAN THORACIC
SOCIETY, AMERICAN MEDICAL ASSOCIATION,
AMERICAN ACADEMY OF PEDIATRICS,
AMERICAN COLLEGE OF PHYSICIANS, AND
LEADERS OF PUBLIC HEALTH SCHOOLS, ET AL.
IN SUPPORT OF RESPONDENTS**

Sara A. Colangelo
Counsel of Record
Jack H.L. Whiteley
GEORGETOWN LAW
ENVIRONMENTAL LAW
& JUSTICE CLINIC
600 New Jersey Ave., NW
Washington, D.C. 20001
(202) 661-6543
sara.colangelo@law.
georgetown.edu

TABLE OF CONTENTS

TABLE OF AUTHORITIES	2
INTEREST OF AMICI.....	3
INTRODUCTION	4
SUMMARY OF ARGUMENT.....	5
ARGUMENT	6
I. Anthropogenic climate change, fueled by emissions of greenhouse gasses such as carbon dioxide, harms public health in the United States.....	6
A. Climate change increases heat-related illnesses, hospitalizations, and death.....	8
B. Climate change fuels longer and more intense fire seasons.....	12
C. Climate change impairs air quality by increasing pollen and ground level ozone.....	15
D. Climate change leads to increased flooding and degraded water quality.....	19
E. Climate change leads to increased vector-borne diseases.....	22
II. Climate change severely harms the health of vulnerable populations.....	25
III. The Clean Air Act empowers EPA to take regulatory action to protect the public from adverse health effects due to climate change.....	30
CONCLUSION.....	33

TABLE OF AUTHORITIES

	Page(s)
Cases	
Placeholder	25
Statutes	
Placeholder	25
Regulatory Materials	
Placeholder	7, 22
Legislative Materials	
Placeholder	19
Other Authorities	
Placeholder.	12

INTEREST OF AMICI¹

Amici are leading physician-member public health organizations, as well as deans, directors, and chairs of public health schools across the country. A complete list of amici and their institutional affiliations appears in Appendix A.

Amici are deeply concerned about the serious and widespread public health harms resulting from anthropogenic climate change, including illness and premature death, and escalating emergency room visits throughout the United States. This brief describes the public health necessity of regulating—and reducing—emissions of greenhouse gases such as carbon dioxide that fuel climate change (collectively, “climate pollutants”). The resolution of this case could have a profound impact on the ability of the Environmental Protection Agency (“EPA” or “the Agency”) to set emissions standards for the largest industrial source of climate pollutants in the United States. Any ruling that reduces the ability of EPA to discharge its public health mission under the Clean Air Act would harm the public welfare. Amici’s collective medical expertise and experience lead them to support the position of the respondents.

¹ Pursuant to Rule 37.6, amici state that no counsel for a party authored this brief in whole or in part and that no person other than amici and their counsel made a monetary contribution to its preparation or submission. All parties filed a blanket consent to the filing of amicus briefs with the exception of the Power Company Respondents. Amici obtained consent from the Power Company Respondents on December 16, 2021.

INTRODUCTION

June 2021 was the warmest month ever recorded on the United States mainland.² Temperatures soared across the Pacific Northwest, reaching 116°F in Portland.³ As record-shattering temperatures seared Oregon and Washington, people flooded emergency rooms seeking care. On June 28th, the U.S. Department of Health and Human Services reported 1,038 emergency room visits for heat-related illness across the Pacific Northwest region; one year earlier on the same day, there were nine.⁴ Hospital systems, already stressed by the COVID-19 pandemic, struggled to admit the surge of patients and medical equipment overheated.⁵

Higher temperatures and punishing heat waves that contribute to illness and injury are two prominent effects of climate change that harm public health. America's leading physician-member medical organizations and public health experts submit this brief to draw the Court's attention to the exigent health threats from climate change. Driven by fossil fuel emissions, climate pollutants harm public health across every segment of

² National Oceanic and Atmospheric Administration, National Centers for Environmental Information, "State of the Climate: Global Climate Report for June 2021," <https://www.ncdc.noaa.gov/sotc/global/202106> (last accessed January 12, 2022).

³ Paul J. Schramm et al., *Heat-Related Emergency Department Visits During the Northwestern Heat Wave – United States, June 2021*, 70(29) *Morbidity and Mortality Weekly Report* 1020 (2021) (noting that temperatures reached 42°F hotter than average June temperatures).

⁴ *Id.*

⁵ Joanne Silberner, *Heat wave causes hundreds of deaths and hospitalizations in Pacific Northwest*, *BMJ* (2021).

American society and in every state. The consequences of climate change impair pulmonary, cardiovascular, neurological, immunological, behavioral health, and other vital systems and functions.

The scale and gravity of these dangers demand regulatory action to reduce emissions of greenhouse gases, including carbon dioxide. Amici urge this Court not to reduce the EPA's ability to regulate carbon dioxide emissions from power plants to protect public welfare and mitigate future public health harms.

SUMMARY OF ARGUMENT

Human-generated greenhouse gas emissions, including carbon dioxide from fossil-fuel combustion, have changed weather patterns and other natural cycles across the world. If left unchecked, this trend will continue, with worsening and compounding public health consequences. In the United States, greenhouse gas-related changes to the weather include more frequent heat waves, higher average temperatures, more forest and urban fires, more air pollution, longer and intensified allergy seasons, more potent and frequent storms and flooding, and expansion in the range of disease-carrying insects. All of these changes will continue to have dangerous health consequences.

These consequences include rises in heat-related illnesses, air pollution-related respiratory and cardiovascular illnesses, injuries and deaths caused by severe fires and storms, the spread of vector-borne diseases like Zika and Dengue, and increases in asthma attack-triggering pollen and mold. The effects of greenhouse gas emissions are occurring in all fifty states, but the harms are not equally distributed. Climate

pollutants' most grievous harms beset children and infants, pregnant women, people over 65, and communities of color and of low income.

Volumes of peer-reviewed science on such health effects reinforce the conclusion that climate pollutants warrant action from EPA. The Clean Air Act authorizes EPA to regulate greenhouse gases as air pollutants as defined under the Act, *Massachusetts v. EPA*, 549 U.S. 497, 528-532 (2007), and mandates their regulation because they endanger public health and welfare. 80 Fed. Reg. 64,510, 64,530 (Oct. 23, 2015). As this Court affirmed, "Congress delegated to EPA the decision whether and how to regulate carbon-dioxide emissions from powerplants." *American Electric Power Co. v. Connecticut*, 564 U.S. 410, 426 (2011).

The EPA's authority to regulate carbon dioxide emissions from power plants is critical to mitigate the scale of health effects of climate pollutants. The Court should be mindful of Congress's decision to provide EPA regulatory authority to address this type of threat to public health. Any retrenchment in the scope of that authority would inflict further harm to the health of current and future generations.

ARGUMENT

I. Anthropogenic climate change, fueled by emissions of greenhouse gasses such as carbon dioxide, harms public health in the United States.

The term "anthropogenic climate change" describes the effects caused by elevated concentrations of greenhouse gases, which trap a higher portion of the

sun's heat that the Earth radiates back into space, leading to rises in global land and ocean temperatures.⁶ In the United States, power plants are “by far” the largest industrial emitters of greenhouse gases. 80 Fed. Reg. at 64,530.

The studies cited in this brief summarize the medical consensus regarding the dire consequences of warming and unstable climate conditions. Report after report establish the escalating toll climate change exacts on public health in the United States. They document the millions of Americans experiencing—and who are predicted to experience—climate change-linked health consequences. Some of these studies measure human health costs in hospitalizations, or missed school and work days.⁷ Others, evaluate them economically in billions of real dollars.⁸ Still others, determine them by quantifying lives shortened and lives lost.⁹

⁶ See, e.g., Environmental Protection Agency, “Causes of Climate Change,” <https://www.epa.gov/climatechange-science/causes-climate-change> (last visited January 12, 2022).

⁷ Neal Fann et al., *The geographic distribution and economic value of climate-change ozone health impacts in the United States in 2030*, 65 *Journal of the Air & Waste Management Association* 570, 574 (2015)

⁸ See, e.g., Kim Knowlton et al., *Six Climate Change-Related Events in the United States Accounted for About \$14 Billion In Lost Lives and Health Costs*, 30 *Health Affairs* 2167 (2011).

⁹ Drew Shindell et al., *The Effects of Heat Exposure on Human Mortality Throughout the United States*, 4 *GeoHealth* 1, 7 (2020) (estimating that during the 2010 decade, 12,000 premature U.S. heat-related deaths occurred annually).

A. Climate change increases heat-related illnesses, hospitalizations, and death.

Climate change results in higher ambient temperatures¹⁰ and more “heat waves,” unusually hot weather that exceeds regional averages for two or more days,¹¹ among other physical transformations. The northern hemisphere is warming faster than the rest of the world, with the northeast suffering the swiftest warming in the contiguous United States.¹² The connection between rising temperatures and health is direct and deadly.¹³ Decades of data from Georgia to

¹⁰ See, e.g., David H. Levinson & Christopher J. Fettig, *Climate Change: Overview of Data Sources, Observed and Predicted Temperature Changes, and Impacts on Public and Environmental Health*, in *Global Climate Change and Public Health* 31, 33–36 (2014) (summarizing leading research on past and projected increases in ambient temperatures).

¹¹ Tiffany T. Smith et al., *Heat waves in the United States: definitions, patterns, and trends*, 118 *Climate Change* 811, 812–13 (2013) (noting that “heat wave” generally refers to temperatures that exceed seasonally and regionally-specific averages for two or more consecutive days); A. Haines et al., *Climate Change and Human Health: Impacts, Vulnerability, and Mitigation*, 367 *Lancet* 2101, 2102 (2006) (concluding that human influence on climate has at least doubled the risk of major heat waves).

¹² Ambarish V. Karmalkar & Raymond S. Bradley, *Consequences of Global Warming of 1.5°C and 2°C for Regional Temperature and Precipitation Changes in the Contiguous United States*, 12 *PloS ONE* e0168697 (2017).

¹³ Shakoor Hajat & Tom Kosatky, *Heat-related mortality: a review and exploration of heterogeneity*, 64 *J. Epidemiology & Cmty. Health* 753 (2010) (determining that risk of mortality in various cities increased by 1–3 percent with each degree-Centigrade increase in temperature); Mercedes Medina-Ramón & Joel Schwartz, *Temperature, temperature extremes, and mortality: a study of*

Washington State demonstrate that intensifying heat resulting from climate change increases emergency room visits for cardiac, pulmonary, and kidney failures, as well as stroke, asthma attacks, and diabetes complications.¹⁴ Even relatively short exposure to extreme heat events is associated with an elevated hourly heart attack rate.¹⁵

Premature heat-related deaths in the continental U.S. are estimated at 12,000 per year.¹⁶ Certain risk factors exacerbate the mortality impacts of heat waves. Large segments of the U.S. population with common pre-existing health conditions, especially the very young or the elderly, are at heightened risk:¹⁷ Multiple studies observe that most heatwave deaths occur in people with cardiovascular or chronic respiratory diseases.¹⁸

acclimatization and effect modification in 50 United States cities, 64 J. Occupational & Env'tl. Med. 827 (2007) (identifying causal relationship based on over six million observations).

¹⁴ Tianqi Chen et al., *Time-series Analysis of Heat Waves and Emergency Department Visits in Atlanta, 1993 to 2012*, 125 Env'tl. Health Perspectives 057009 (2017); Tania Busch Isaksen et al., *Increased hospital admissions associated with extreme-heat exposure in King County, Washington, 1990-2010*, Reviews on Environmental Health 10.1515/reveh-2014-0050 (2015).

¹⁵ Sebastian T. Rowland et al., *Can ultra short-term changes in ambient temperature trigger myocardial infarction?*, 143 Environment International, 105910 (2020).

¹⁶ See, e.g., Shindell et al., *Effects of Heat*, *supra* note __, at 7.

¹⁷ See, e.g., Rupa Basu, *High ambient temperature and mortality: a review of epidemiologic studies from 2001 to 2008*, 8 Env'tl. Health, __ (2009) (determining that the groups most vulnerable to elevated heat-related deaths included infants and young children, and those over 65).

¹⁸ Anthony J. McMichael et al., *Climate change and human health: present and future risks*, 367 Lancet 9513, at 861 (2006).

Residents in urban areas also suffer from the “heat island” effect of concrete surfaces heating faster and holding heat longer than vegetation and water surfaces prevalent in non-urban areas.¹⁹ And populations living in locations with historically lower temperatures often lack air conditioning and other adaptations, and thus experience higher mortality rates from heat waves.²⁰

Heat waves and higher temperatures also cause a number of other serious health effects. One effect is “heat stress,” when the body receives heat “in excess of what it can tolerate without physiological impairment.”²¹ U.S. labor productivity impairment from heat stress is projected to double between 1995 and 2030.²² “The expected productivity loss in 2030 is equivalent to 389,000 full-time jobs... concentrated in the southern states... and concern[ing] mostly outdoor workers, such as construction workers and farm[ers]... .”²³

¹⁹ Clare Heaviside et al., *The Urban Heat Island: Implications for Health in a Changing Environment*, 4(3) Current Environmental Health Reports 296 (2017).

²⁰ William N. Rom & Kent E. Pinkerton, *Introduction: Consequences of Global Warming to the Public’s Health*, in *Global Climate Change and Public Health* 1, 10 (2014).

²¹ Tord Kjellstrom et al., *Heat, human performance, and occupational health: A key issue for the assessment of global climate change impacts*, 37 *Annual Review of Public Health* 97, 98 (2016).

²² International Labour Organization, *Working on a Warmer Planet: The Impact of Heat Stress on Productivity and Decent Work*, at 43 (2019).

²³ *Id.*

Other health harms associated with heat waves and higher temperatures include heat stroke,²⁴ adverse birth outcomes for pregnant women,²⁵ and decreased lung function.²⁶ For example, a study of 12.5 million Medicare beneficiaries across 213 U.S. counties found that each 10°F increase in daily temperature was associated with a 4.3 percent increase in same-day emergency hospitalizations for respiratory diseases.²⁷ Further, extreme heat has significant adverse effects on mental health.²⁸ Heat waves impair cognition, moods, and

²⁴ R. Sari Kovats & Shakoor Hajat, *Heat stress and public health: a critical review*, 29 Annual Rev. Pub. Health 41, 42, 47 (2008) (noting danger of and risk factors for heat stroke); Helene G. Margolis, *Heat Waves and Rising Temperatures: Human Health Impacts and the Determinants of Vulnerability*, in *Global Climate Change and Public Health*, 85, 97-100 (2014) (describing pathways through which high temperatures can lead to adverse health outcomes).

²⁵ Bruce Bekkar et al., *Association of Air Pollution and Heat Exposure With Preterm Birth, Low Birth Weight, and Stillbirth in the US: A Systematic Review*, 3 JAMA Network Open e208243 (2020) (providing a review of 57 studies and concluding heat, ozone, and fine particulate matter are all associated with preterm birth, low birth weight, and stillbirth).

²⁶ Mary B. Rice et al., *Association of outdoor temperature with lung function in a temperate climate*, 53(1) Eur Respir J. 1, 1 (2019) (establishing that “1-, 2- and 7-day [higher] average temperatures were all associated with lower lung function.”); see also Nana Mireku et al. *Changes in weather and the effects on pediatric asthma exacerbations*, 103 Annals of Allergy, Asthma & Immunology 220 (2009).

²⁷ See G. Brooke Anderson et al. *Heat-related emergency hospitalizations for respiratory diseases in the Medicare population*, 187 Am. J. Respir. Crit. Care Med. 1098 (2013).

²⁸ Nick Obradovich et al., *Empirical evidence of mental health risks posed by climate change*, 115 Proc. Nat’l Acad. Sci. 10953 (2018); Jingwen Liu et al., *Is there an association between hot weather and poor*

sleep,²⁹ and contribute to increases of aggression and suicide.³⁰ Without curbing greenhouse gas emissions, ambient temperatures and heat waves will intensify, with profound consequences for human health.³¹

B. Climate change fuels longer and more intense fire seasons.

Wildfires and fires in densely populated areas lead directly to loss of life and property, and are increasing in frequency, duration, and intensity.³² Multiple studies

mental health outcomes? A systematic review and meta-analysis, 153 *Environment Int'l* 106533 (2021).

²⁹ R. Jisung Park et al., *Learning is inhibited by heat exposure, both internationally and within the United States*, *Nature Human Behavior* (2020); N. Obradovich et al., *Nighttime temperature and human sleep loss in a changing climate*, 3 *Science Advances*. E1601555 (2017).

³⁰ Marshall Burke et al., *Climate and Conflict*, 7 *Annual Rev. of Econ.* 577 (2015); Marshall Burke et al., *Higher Temperatures increase suicide rates in the United States and Mexico*, 8 *Nature Climate Change* 723 (2018).

³¹ Roger D. Peng et al., *Toward a Quantitative Estimate of Future Heat Wave Mortality under Global Climate Change*, 119 *Envtl. Health Persp.* 701, 701 (2011) (“The impact of future heat waves on human health will likely be profound, and significant gains can be expected by lowering future carbon dioxide emissions.”).

³² See, e.g., Philip E. Dennison et al., *Large wildfire trends in the western United States, 1984-2011*, 41 *Geophys. Res. Letters* 2928 (2014) (number of large U.S. fires increasing); Jeremy S. Littell et al., *Climate and wildfire area burned in western U.S. ecoregions*, 19(4) *Ecological Applications* 1003 (2009) (U.S. area burned in fires increasing); A.L. Westerling et al., *Warming and earlier spring increase western U.S. forest wildfire activity*, 313 *Science* 940 (2006) (U.S. fire season duration increasing).

conclude that worsening fire seasons are largely attributable to climate change, due to increases in temperatures and aridity, and earlier snowmelt.³³ The United States became significantly more wildfire-prone in the past four decades,³⁴ with the area burned from wildfires quadrupling.³⁵ Wildfires release high concentrations of pollutants, including particulate matter and chemical compounds which form ground-level ozone.³⁶ These pollutants harm populations far from the western continental United States where fires

³³ See, e.g., Mary B. Rice et al., *Respiratory Impacts of Wildland Fire Smoke: Future Challenges and Policy Opportunities*, 18 *Annals of the American Thoracic Society* 921 (2021) (observing wildfire activity increases are largely attributable to climate change rather than land use or forest management); John T. Abatzoglou & A. Park Williams, *Impact of anthropogenic climate change on wildfire across western US forests*, 42 *Proceedings of the National Academy of Sciences* 113 (2016) (“human-caused climate change caused over half of the documented increases in fuel aridity since the 1970s and doubled the cumulative forest fire area since 1984”).

³⁴ See Dennison et al., *Large Wildfire Trends*, *supra* note 1, at 2932-33; Steven W. Running, *Is Climate Change Causing More, Larger Wildfires?*, 313 *Science* 927, 927 (2006) (reporting a fourfold increase in major American wildfires since 1986).

³⁵ Marshall Burke et al., *The Changing Risk and Burden of Wildfire in the United States*, 118 *Proc. Nat’l Academy of Science* 1, 1 (2021).

³⁶ Jennifer D. Stowell et al., *Associations of wildfire smoke PM2.5 exposure with cardiorespiratory events in Colorado 2011-2014*, 133 *Environment International* 105151 (2019) (demonstrating that increased exposure to wildfire-derived PM2.5 was associated with increased respiratory hospitalizations, when separating out background PM); Daniel A. Jaffe & Nicole L Wigder, *Ozone production from wildfires: A critical review*, 51 *Atmospheric Env’t* 1, 2, 7 (2012).

typically burn, reaching Midwestern and Northeastern states.³⁷ Indeed, as of 2017, an estimated 10 percent of the U.S. population, approximately 30.5 million people, reside where wildfire can contribute a significant burden to their exposure to fine particulate matter.³⁸

Wildfire-generated particulate matter increases respiratory and cardiac hospitalizations. For example, studies in Colorado and California demonstrate marked escalation in emergency visits and hospitalizations for asthma from wildfire-generated particulate matter.³⁹ There is also strong evidence that exposure to particulate matter increases risk of death, even for those without preexisting conditions.⁴⁰ Further, the growing severity and frequency of weather-related climate disasters harm mental health, as well, when wildfires and storms destroy homes and communities.⁴¹ Responses range

³⁷ See, e.g., Katelyn O'dell et al., *Estimated Mortality and Morbidity Attributable to Smoke Plumes in the United States: Not Just a Western US Problem*, 5(9) *GeoHealth* e2021GH000457 (2021).

³⁸ A.G. Rappold et al., *Community Vulnerability to Health Impacts of Wildland Fire Smoke Exposure*, 51 *Environmental Science and Technology*, 6674 (2017).

³⁹ Stowell et al., *Associations of wildfire smoke PM2.5*, *supra* note

—
⁴⁰ Ana G. Rappold et al., *Cardio-respiratory outcomes associated with exposure to wildfire smoke are modified by measures of community health*, 11 *Envtl. Health*, Sept. 24, 2012; Johanna Lepeule et al., *Chronic Exposure to Fine Particles and Mortality: An Extended Follow-up of the Harvard Six Cities Study from 1974 to 2009*, 120 *Envtl. Health Persps.* 965, 968 (2012).

⁴¹ Ozone and particulate matter exposure is linked to increased incidence of depression, anxiety, and dementia. See Isobel Braithwaite et al., *Air Pollution (Particulate Matter) Exposure and Associations with Depression, Anxiety, Bipolar, Psychosis and Suicide*

from post-traumatic stress disorder, to new onset or exacerbation of psychiatric disorders, to complex grief, among other disorders.⁴²

C. Climate change impairs air quality by increasing pollen and ground level ozone.

1. Pollen

Climate change is the dominant driver of the United States' lengthening pollen season and a significant contributor to increasing pollen concentrations.⁴³ Warmer temperatures lengthen pollen seasons because plants bloom earlier and for longer periods of time.⁴⁴ In addition, climate change's meteorological effects include more frequent and severe thunderstorms, which cause

Risk: A Systematic Review and Meta-Analysis, 127 *Envtl. Health Persp.* 126002 (2019).

⁴² Hayes et al. *Climate change and mental health: risks, impacts and priority actions*, 12 *Intern. J. of Mental Health Sys* (2018).

⁴³ See, e.g., William R.L. Anderegg, *Anthropogenic Climate Change Is Worsening North American Pollen Seasons*, 118 *Proc. Nat'l Academy of Science* 1 (2021); L.H. Ziska & D. Berman, *Impact of Climate Change on Aeroallergenic Pollen Metrics: A Hemispheric Perspective*, 33 *Current Allergy & Clinical Immunology* 93 (2020); Yong Zhang et al., *Allergic pollen season variations in the past two decades under changing climate in the United States*, 21 *Global Change Biology* 1581, 1583-86 (2015).

⁴⁴ Lewis Ziska et al., *Recent Warming by Latitude Associated with Increased Length of Ragweed Pollen Season in Central North America*, 108 *Proc. Nat'l Academy of Science* 4248, 4248 (2011) (documenting that between 1995 and 2009, the ragweed pollen season lengthened by 13-27 days above the forty-fourth parallel, which encompasses portions of the United States).

sudden pollen releases⁴⁵ and also break pollen into smaller particles, enabling its allergens to penetrate deeper into the lungs.⁴⁶ The predictable result: more asthma attacks and more emergency room visits.⁴⁷

Like the heat-related dangers described above, the impacts of pollen are more severe for people with pre-existing health conditions. Longer and more intense allergy seasons pose a substantial threat to the approximately 25.1 million Americans with asthma,⁴⁸ because pollen triggers attacks in asthmatics who are allergic to pollen.⁴⁹ Nearly 9 percent of the nation's school age children have asthma.⁵⁰ And asthma exacerbations keep children out of school and adults out

⁴⁵ Shuaib M. Nasser & Thomas B. Pulimood, *Allergens and Thunderstorm Asthma*, 9 *Current Allergy & Asthma Rep.* 384, 387-88 (2009).

⁴⁶ Andrew Rorie & Jill A. Poole, *The Role of Extreme Weather and Climate-Related Events on Asthma Outcomes*, 41(1) *Immunology & Allergy Clin. N. Am.* 73 (2021).

⁴⁷ James E. Neumann et al., *Estimates of Present and Future Asthma Emergency Department Visits Associated with Exposure to Oak, Birch, and Grass Pollen in the United States*, 10 *GeoHealth* 11 (2019) (determining health impacts for exposure to current and potential future pollen loads under multiple climate scenarios).

⁴⁸ Centers for Disease Control and Prevention, *Most Recent National Asthma Data*, https://www.cdc.gov/asthma/most_recent_national_asthma_data.htm (last accessed January 18, 2022).

⁴⁹ Clarisse Gautier & Denis Charpin, *Environmental triggers and avoidance in the management of asthma*, 10 *J. of Asthma and Allergy*, 47 (2017); Susan C. Anenberg et al., *Impacts of Oak Pollen on Allergic Asthma in the United States and Potential Influence of Future Climate Change*, 1 *GeoHealth* 80, 90 (2009).

⁵⁰ Centers for Disease Control and Prevention, *supra* _ (calculating the number of children 5-17 in the U.S. with asthma).

of work.⁵¹ Recurrent exacerbations can cause permanent airway damage and often require costly medical care.⁵²

2. Ground-Level Ozone

Warmer temperatures that come with higher atmospheric concentrations of greenhouse gases increase ground-level ozone.⁵³ Ground-level ozone is created through a photochemical reaction between nitrogen oxides, volatile organic compounds, heat and sunlight.⁵⁴ It causes difficulty breathing, coughing and shortness of breath, and contributes to respiratory-related death.⁵⁵

People suffering from pre-existing lung disease, and children with still-developing lungs, are especially

⁵¹ Susan M. Pollart et al., *Management of acute asthma exacerbations*, 84 *Am. Family Physician* 40 (2011); Mary E. Streck, *Difficult asthma*, 3 *Proc. Am. Thoracic Soc'y* 116 (2006).

⁵² Gary S. Rachelefsky, *From the page to the clinic: Implementing new National Asthma Education and Prevention Program guidelines*, 9 *Clinical Cornerstone* 9, 9–10 (2009).

⁵³ Fann et al., *The geographic distribution*, *supra* note __, at 570 (“Climate change can affect air pollutant concentrations in a myriad of ways. Meteorological factors, such as temperatures, cloudiness, precipitation frequency and intensity, ... all... influence air quality by ... determining photochemical reaction rates....”); Ivar Isaksen et al., *Atmospheric composition change: Climate-Chemistry interactions*, 43 *Atmospheric Environment* 5138 (2009).

⁵⁴ Fann et al., *Geographic Distribution*, *supra* note __, at 570.

⁵⁵A. Wilson et al., *Climate change impacts on projections of excess mortality at 2030 using spatially varying ozone–temperature risk surfaces*. *J Expo Sci Environ Epidemiol* **27**, 118–124 (2017) (modeling ozone-related mortality due to projected changes in climate conditions); Jennifer D. Stowell et al., *The impact of climate change and emissions control on future ozone levels: Implications for human health*, *Environment International* (2017) (discussing health benefits of emissions mitigation).

susceptible to the harmful effects of ozone exposure. Even modest and relatively brief increases in ground-level ozone are linked to an elevated risk of hospitalization for patients with asthma and chronic obstructive pulmonary disease.⁵⁶ Tens of thousands of additional ozone-related premature deaths and illnesses are predicted to occur by 2030 if emission controls are not adopted.⁵⁷

D. Climate change leads to increased flooding and degraded water quality.

Climate change-linked higher temperatures lead to more extreme rainfall over short periods of time, producing dangerous floods.⁵⁸ “[W]armer air is capable of holding more water than cooler air, and therefore has the potential to provide more moisture to rainfall events.”⁵⁹ Extreme floods in the United States have increased by more than twenty percent in recent decades

⁵⁶ See Benedicte Jacquemin et al., *Air pollution and asthma control in the Epidemiological study on the Genetics and Environment of Asthma*, 66 J. Epidemiology Community Health 796-802 (2012); Kelly Moore, et al., *Ambient ozone concentrations cause increased hospitalizations for asthma in children: an 18-year study in Southern California*, 116 *Envtl. Health Perspectives* 1063-70 (2008).

⁵⁷ Fann et al., *Geographic distribution*, *supra* note 56; see also Kim Knowlton et al., *Assessing Ozone-Related Health Impacts under a Changing Climate*, 112 *Envtl. Health Persps.* 1557, 1559-60, 1562 (2004) (estimating significant increase in mortality by 2050 as a result of increase in ground-level ozone attendant to climate change).

⁵⁸ Seth Westra et al., *Future Changes to the Intensity and Frequency of Short-Duration Extreme Rainfall*, 52 *Rev. of Geophysics* 522, 522-25 (2014).

⁵⁹ *Id.*

in some regions.⁶⁰ For example, “heat stress events,” in which an environment is struck by high temperatures and humidity, preceded a high percentage of recent floods in Iowa, Illinois, and Indiana.⁶¹

The potential effects of heat stress-linked floods include a high number of fatalities and the destruction of communities and critical infrastructure.⁶² For example, climate-related flooding can damage roads, hospitals, and the nation’s power grid.⁶³ The health effects and costs are likewise significant. For instance, a 2011 study measured the health costs of river flooding.⁶⁴ Floods in North Dakota from severe storms and near-record snow accumulation and rapid melting, exacted approximately \$145,495 in health costs per 1,000 people in the area affected.⁶⁵

The cascading impacts of climate change-fueled flooding include illnesses from microbial growth. After

⁶⁰ Wouter R. Berghuijs et al., *Recent Changes in Extreme Floods Across Multiple Continents*, 12 *Envtl. Res. Letters* 1, 4 (2017) (estimating increases in the occurrence of extreme floods by region throughout the world).

⁶¹ See Wei Zhang & Gabriele Villarini, *Deadly Compound Heat Stress-Flooding Hazard Across the Central United States*, 15 *Geophysical Research Letters* 1, 6 (2020) (“There is a clear connection between heat stress and flooding.... The new compounding extreme exhibits a strong signal, especially in Iowa, Illinois, and Indiana, which are frequently affected by stormy weather during the summer.”).

⁶² *Id.* at 6.

⁶³ *Id.*

⁶⁴ Knowlton et al., *Six Climate Change-Related Events*, *supra* note

⁶⁵ *Id.* at 2169-70.

Hurricanes Katrina and Rita made landfall in 2005, water inundated 80 percent of New Orleans for more than two weeks.⁶⁶ Even as floodwaters receded, additional threats emerged. The flooding's duration, coupled with warm temperatures in the late Louisiana summer, spawned mold in thousands of homes.⁶⁷ Exposure to mold is associated with respiratory illnesses, placing immunocompromised people at risk for fungal colonization and opportunistic infections.⁶⁸ For example, following floods in North Dakota and North Carolina, public health workers reported mold-linked, post-flooding increases in asthma symptoms, rhinitis, rash, and headaches.⁶⁹

Extreme flooding also diminishes water quality. Increased rain and snow can exceed the capacity of sewer

⁶⁶ See Margaret A. Riggs et al., *Resident Cleanup Activities, Characteristics of Flood-Damaged Homes and Airborne Microbial Concentrations in New Orleans, Louisiana, October 2005*, 106 *Env'tl. Research* 401, 402 (2005). Climate change also increases the intensity of hurricanes. See Greg Holland & Cindy L. Bruyère, *Recent Intense Hurricane Response to Global Climate Change*, 42 *Climate Dynamics* 617, 617-19 (2013).

⁶⁷ *Id.* at 402, 404-07.

⁶⁸ *Id.* (citing Institute of Medicine, *Damp Indoor Spaces and Health* (2004)). After the floods, families enrolled in a study conducted by a Louisiana asthma association reported moving homes between one and four times to find safe housing. Jill A. Poole et al., *Impact of Weather and Climate Change with Indoor and Outdoor Air Quality in Asthma: A Work Group Report of the AAAAI Environmental Exposure and Respiratory Health Committee*, 143 *J. Allergy Clinical Immunology* 1702, 1705 (2019).

⁶⁹ Riggs et al., *Resident Cleanup Activities*, *supra* note __, at 402.

systems.⁷⁰ Discharges from domestic, commercial, and industrial sources of waste can then flow directly into surface waters, including rivers, streams, and estuaries.⁷¹ These discharges in turn impair water quality, expose people to untreated sewage, and lead to basement backups of sewage in residential homes.⁷² These exposures cause gastrointestinal illness and other waterborne diseases.⁷³ River flooding further degrades water quality by raising the rates of dissolved nitrogen, phosphorus, and suspended solids, as well as overwhelming wastewater treatment plants.⁷⁴ Ample data indicate that as climate change-linked flooding worsens, its human health costs are likely to multiply.⁷⁵

E. Climate change leads to increased vector-borne diseases.

Vector-borne diseases result from infections transmitted by mosquitoes and ticks. The expanding range of both mosquitoes and ticks, and the pathogens they carry, is attributable to an array of human-induced changes, including climate change. Temperatures are

⁷⁰ U.S. Environmental Protection Agency, Report to Congress: Combined Sewer Overflows into the Great Lakes Basin 1 (2016).

⁷¹ *Id.* at 1-2.

⁷² *Id.* at 2.

⁷³ See, e.g., Jonathan A. Patz et al., *Climate Change and Waterborne Disease Risk in the Great Lakes Region of the U.S.*, 35 Am. J. Preventive Med. 451, 455 (2008).

⁷⁴ Thomas C. Peterson et al., *Changes in Weather and Climate Extremes: State of Knowledge Relevant to Air and Water Quality in the United States*, 64 J. of the Air & Waste Management Assoc. 184, 191 (2014).

⁷⁵ Stephane Hallegatte et al., *Future Flood Losses in Major Coastal Cities*, 3 Nature Climate Change 802, 804-05 (2013).

central to mosquito physiology and mortality, to their host behavior, and to the incubation of pathogens within the mosquito.⁷⁶ Warmer weather thus enables mosquitoes to expand their range.⁷⁷

Over the last several decades, the expanded range of multiple mosquito species facilitated the spread of serious vector-borne diseases into the United States. Physicians attribute the recent proliferation of mosquito-borne illnesses such as Zika—which causes fetal neurological complications and birth defects including microcephaly—to rising global surface temperatures and new variability in rainfall.⁷⁸ In fact, although Zika became transmissible to humans around 1950, the United States had no reported local transmissions until 2016.⁷⁹

Dengue is another mosquito-borne illness with a recently-expanded range.⁸⁰ Previously limited to subtropical and tropical regions, Dengue outbreaks now

⁷⁶ Michael A. Robert et al., *Climate Change and Viral Emergence: Evidence from Aedes-Borne Arboviruses*, 40 *Current Opinion in Virology* 41, 42 (2020).

⁷⁷ See Ilia Rochlin et al., *Climate Change and Range Expansion of the Asian Tiger Mosquito (Aedes Albopictus) in Northeastern USA: Implications for Public Health Practitioners*, 8 *PLoS One* 1, 1-2 (2013). Climate change-influenced extreme weather events also produce conditions in which water-, mosquito-, and rodent-borne diseases can thrive. See, e.g., Paul Epstein, *The Ecology of Climate Change and Infectious Diseases: Comment*, 91 *Ecology* 925 (2010).

⁷⁸ Robert et al., *Viral Emergence*, *supra* note , at 41-44.

⁷⁹ Nathan D. Grubaugh et al., *Genomic Epidemiology Reveals Multiple Introductions of Zika Virus into the United States*, 546 *Nature* 401, 401-02 (2017).

⁸⁰ Robert et al., *Viral Emergence*, *supra* note , at 42. (observing an expansion of Dengue in the last 20 years).

occur in Hawaii, Florida, and Texas.⁸¹ Strikingly, the range of the Dengue-carrying mosquito has now grown to include the entire southeast and much of the southwest United States.⁸²

The introduction of the West Nile Virus into the United States is also linked to climate, particularly the proliferation of warm and wet conditions.⁸³ It is now the most prevalent mosquito-borne disease in the United States.⁸⁴ West Nile Virus can attack the central nervous system, necessitating hospitalization, and sometimes causing death.⁸⁵

Mosquitoes are not the only disease-inducing pest with an enlarged range. The habitats of several tick species are growing throughout the United States in response to increased temperatures.⁸⁶ This warming trend contributes to tick species' decades-long encroachment to the north and west.⁸⁷ Ticks cause almost 95 percent of all vector-borne diseases reported annually

⁸¹ *Id.* at 42.

⁸² Centers for Disease Control and Prevention, *Potential Range of the Aedes aegypti and Aedes albopictus in the United States, 2017*, available at <https://www.cdc.gov/mosquitoes/mosquito-control/professionals/range.html> (accessed January 2, 2022).

⁸³ Epstein, *Ecology of Climate Change*, *supra* note __, at 927.

⁸⁴ Carolyn A. Reimann et al., *Epidemiology of Neuroinvasive Arboviral Disease in the United States, 1999-2007*, 79 *Am. J. Trop. Med. Hyg.* 974, 974 (2008).

⁸⁵ *Id.*

⁸⁶ Daniel E. Sonenshine, *Range Expansion of Tick Disease Vectors in North America: Implications for Spread of Tick-Borne Disease*, 15 *Int'l J. Env'tl. Res. Public Health*, 478 (2018).

⁸⁷ *Id.*

in the United States, including Lyme disease and Rocky Mountain spotted fever, among others.⁸⁸

In short, as temperatures rise, the range of environments suitable for disease-carrying species grows.⁸⁹ In the absence of effective regulation, regions affected by vector-borne illnesses are likely to expand, new vector-borne diseases may emerge, and existing vector-borne diseases may increase.⁹⁰

II. Climate change severely harms the health of vulnerable populations.

The health harms and costs of climate pollutants fall heavily on vulnerable populations. Young children and pregnant women, adults older than sixty-five, and communities of color and low income are most vulnerable to and severely harmed by the adverse health impacts of climate change.⁹¹

Children, particularly infants, are more susceptible to climate change-related temperature increases and heat waves because they cannot regulate body temperature as well as adults.⁹² Children are also at heightened risk from

⁸⁸ *Id.*

⁸⁹ See Rochlin et al., *Asian Tiger Mosquito*, *supra* note __, at 1-2.

⁹⁰ Ahdoot & Pacheco, *supra* note __, at e1474.

⁹¹ See, e.g., Yunquan Zhang et al., *Socio-geographic disparity in cardiorespiratory mortality burden attributable to ambient temperature in the United States*, 26 *Envtl. Science and Pollution Research Int'l* 694 (2019).

⁹² Kim Knowlton et al., *The 2006 California heat wave: Impacts on hospitalizations and emergency department visits*, 117 *Envtl. Health Persps.* 61 (2009) (observing greater risk of heat-related emergency department visits for children ages 0-4); Aaron S. Bernstein &

air pollution harms because they generally spend more time outdoors, have higher respiratory rates, and have developing organs and immune systems.⁹³ Exposure of children to air pollutants is associated with reduced lung function, new onset asthma, exacerbation of chronic respiratory illnesses,⁹⁴ cognitive and developmental disorders including autism and attention deficit/hyperactivity disorder,⁹⁵ and asthma-related hospitalizations.⁹⁶

The risk of severe heat complications for pregnant women and infants also escalates with temperature increases.⁹⁷ One 2019 study based on 32 million births across 403 U.S. counties established that exposure to extreme heat is associated with an increased risk of

Samuel S. Myers, *Climate Change and Children's Health*, 23 *Current Opinion in Pediatrics* 221, 222 (2011).

⁹³ Heather Brumber et al., *American Academy of Pediatrics, Ambient Air Pollution: Health Hazards to Children*, 147 *Pediatrics* e2021051484 (2021).

⁹⁴ Brumber, *supra* note 93; Gennaro D'Amato et al., *Urban Air Pollution and Climate Change as Environmental Risk Factors of Respiratory Allergy: An Update*, 20 *J. Investigational Allergology & Clinical Immunology* 95 (2010).

⁹⁵ Oddvar Myhre et al., *Early life exposure to air pollution particulate matter (PM) as risk factor for attention deficit/hyperactivity disorder (ADHD): Need for novel strategies for mechanisms and causalities*, 354 *Toxicol. Appl. Pharmacol.* 196 (2018); Andrea L. Roberts et al., *Perinatal Air Pollutant Exposures and Autism Spectrum Disorder in the Children of Nurses' Health Study II participants*, 3 *Env'tl. Health Persps.* 264 (2013).

⁹⁶ See, e.g., Mireku et al., *Asthma exacerbations*, *supra* note 93, at 223-24; Katherine Shea, *Global Climate Change and Children's Health*, 120 *Pediatrics* 1359 (2007).

⁹⁷ Bekkar et al., *Association of Air*, *supra* note 93.

preterm birth, the second leading cause of death in children under five.⁹⁸ Further, heat waves during pregnancy are correlated with increased maternal stress and, consequently, babies with abnormal conditions related to maternal stress.⁹⁹ In addition, higher incidences of wildfires are dangerous for pregnant women, and infants with developing lungs and brains.¹⁰⁰ Wildfire smoke exposure during pregnancy is associated with low birth weight and pre-term birth.¹⁰¹

Climate change also presents more serious health threats to people 65 and older.¹⁰² People over 65 are more likely to have a pre-existing condition that renders climate co-morbidity more likely. For example, older

⁹⁸ Shengzhi Sun et al., *Ambient temperature and preterm birth: A retrospective study of 32 million US singleton births*, 126 *Environment International* 7 (2019).

⁹⁹ Gulcan Cil & Trudy Anne Cameron, *Potential Climate Change Health Risks from Increases in Heat Waves: Abnormal Birth Outcomes and Adverse Maternal Health Conditions*, 37 *Risk Analysis* 2066 (2017) (examining adverse conditions such as fetal distress and reliance on a ventilator at birth).

¹⁰⁰ See, e.g., Sam Heft-Neal et al., *Associations between wildfire smoke exposure during pregnancy and risk of preterm birth in California*, 203 *Environmental Research* 111872 (2022).

¹⁰¹ Sana Amjad et al., *Wildfire exposure during pregnancy and the risk of adverse birth outcomes: A systematic review*, 156 *Environment Int'l* 106644 (2021).

¹⁰² Rice et al., *Respiratory Impacts of Wildland Fire*, *supra* note __, at 923; Carina J. Gronlund et al., *Vulnerability to Renal, Heat and Respiratory Hospitalizations During Extreme Heat Among U.S. Elderly*, 136(3) *Climatic Change*, 631 (2016). See also Gill Livingston et al., *Dementia prevention, intervention and care: 2020 report of the Lancet Commission*, 396 *Lancet Commissions* 413 (2020) (describing the risk of dementia from exposure to pollutants associated with climate change).

people are more likely to be hospitalized or to die from high temperatures and heat waves.¹⁰³ This population has marginal cardio-respiratory reserves to cope with heat and air pollution, placing them at risk for more frequent acute cardiovascular and respiratory illnesses.¹⁰⁴

But age is not the only predictor of climate change health effects. Race and income are significant predictors of these risks.¹⁰⁵ In a study tracking more than a decade of heat-related adult deaths in New York City, researchers found that Black adults and those living in census tracts receiving greater public assistance were most likely to die during heat waves.¹⁰⁶ In the western wildfire context, studies also establish that Black adults and children are at significantly higher risk of

¹⁰³ Ambarish Vaidyanathan et al., *Heat-Related Deaths – United States, 2004-2018*, 69 *Morbidity and Mortality Weekly Report* 729 (2020); Janet L. Gamble et al., *Climate Change and Older Americans: State of the Science*, 121 *Envtl. Health Persps.* 15 (2013).

¹⁰⁴ Antonella Zanobetti et al., *Summer temperature variability and long-term survival among elderly people with chronic disease*, 109 *Proc. Nat'l Acad. Sci.* 6608 (2012); Anderson et al., *Heat-Related Emergency*, *supra* note _ at 1098.

¹⁰⁵ See, e.g., Zhang et al., *Socio-geographic disparity*, *supra* note _.

¹⁰⁶ Jaime Madrigano et al., *A Case-Only Study of Vulnerability to Heat Wave-Related Mortality in New York City (2000-2011)*, 123 *Envtl. Health Persps.* 672, 672 (2015) (“Compared with other warm-season days, deaths during heat waves were more likely to occur in black (non-Hispanic) individuals than other race/ethnicities, . . . and more likely among those living in census tracts that received greater public assistance....”).

respiratory-related hospital admissions on high smoke days.¹⁰⁷

Communities of color are not just at a heightened risk of health impacts from climate pollutants. Often they experience a disproportionate burden of multiple health stressors. The disparities in exposure to air pollution at home and at school by race and ethnicity is documented in several studies.¹⁰⁸ As a result, residents in these communities suffer elevated rates of conditions that render them more vulnerable to health harms from climate change, such as asthma, chronic airway diseases, and cardiovascular disease.¹⁰⁹ These communities, then, along with children, infants, pregnant women, and those over 65, are at the greatest risk if greenhouse gas emissions are not regulated and reduced.

III. The Clean Air Act empowers EPA to take regulatory action to protect the public from adverse health effects due to climate change.

The purpose of the Clean Air Act is “to protect and enhance the quality of the Nation’s air resources so as to

¹⁰⁷ Jia Coco Liu et al., *Who Among the Elderly Is Most Vulnerable to Exposure to and Health Risks of Fine Particulate Matter From Wildfire Smoke?*, 186 *American J. of Epidemiology* 730 (2017).

¹⁰⁸ See, e.g., Jonathan Colmer et al., *Disparities in PM2.5 air pollution in the United States*, 369 *Science* 575 (2020); Jayajit Chakraborty et al., *Children at Risk: Measuring Racial/Ethnic Disparities in Potential Exposure to Air Pollution at School and Home*, 61 *J. Epidemiology & Cmty. Health* 1074 (2017); Eric B. Brandt et al., *Air Pollution, Racial Disparities, and COVID-19 Mortality*, 146 *J. Allergy & Clinical Immunology* 61, 62 (May 7, 2020) (“Lower income communities of color are more likely to have historical exposures to higher levels of air pollution.”).

¹⁰⁹ Eric B. Brandt et al., *supra* note _.

promote the public health and welfare... ." 42 U.S.C. § 7401(b)(1). Public health was of paramount importance to Congress in drafting and amending the CAA. Senator Edmund Muskie, the Senate architect of the legislation, said the Act would "protect the public health," noting that the costs of air pollution included "death, disease, and disability." 116 CONG. REC. S20,597 (1970).¹¹⁰ The text of the CAA is replete with mandates for EPA to consider public health in its decision-making and to protect public health with its actions. *See* 42 U.S.C. §§ 7408-09 (air quality criteria and national ambient air quality standards); § 7411 (standards of performance for new stationary sources); § 7412 (hazardous air pollutants); § 7470 (prevention of significant deterioration); § 7521 (emission standards for new motor vehicles). This Court's decisions also reflect the Act's textual commitment to public health. *See, e.g., Whitman v. American Trucking Ass'n*, 531 U.S. 457, 465-71 (2001).

Accordingly, from the earliest days of CAA implementation, EPA regulated power plants because they pose an array of risks to public health. *See, e.g.,* 36 Fed. Reg. 5931 (Mar. 31, 1971) (memorializing EPA's decision in 1971 that coal-fired generators fit under sources to be regulated by Section 111 because they "[contribute] significantly to the endangerment of public health or welfare.") And more than a decade ago, EPA concluded that greenhouse gases from automobiles, also emitted from power plants, threatened the public health and welfare of current and future generations. U.S. EPA, *Endangerment and Cause or Contribute Findings for*

¹¹⁰ In introducing the Senate legislation for the 1990 CAA amendments, Senator Lincoln Chafee emphasized, "[t]his is a health bill." 136 CONG. REC. S16895.

Greenhouse Gases Under Section 202(a) of the Clean Air Act, 74 Fed. Reg. 66,496 (Dec. 15, 2009).

Regulation and reduction of greenhouse gas emissions are necessary to mitigate the scale of intensifying public health harms associated with climate change. The need is urgent, and the quality and length of lives are at stake. The Court should affirm EPA's ability to carry out its mandate to protect public health by regulating carbon dioxide emissions from power plants.

CONCLUSION

The judgment of the court of appeals should be affirmed.

Respectfully submitted,

Sara A. Colangelo

Counsel of Record

Jack H.L. Whiteley

GEORGETOWN LAW

ENVIRONMENTAL LAW

& JUSTICE CLINIC

600 New Jersey Ave., NW

Washington, D.C. 20001

(202) 661-6543

sara.colangelo@law.

georgetown.edu

Counsel for Amici Curiae

January 25, 2022